

The Shannon information of filtrations and the additional logarithmic utility of insiders

Stefan Ankirchner
Institut für Mathematik
Humboldt-Universität zu Berlin
Unter den Linden 6
10099 Berlin
Germany

Steffen Dereich
Fachbereich Mathematik
Technische Universität Berlin
Strasse des 17. Juni 136
10623 Berlin
Germany

Peter Imkeller
Institut für Mathematik
Humboldt-Universität zu Berlin
Unter den Linden 6
10099 Berlin
Germany

July 15, 2004

Abstract

The background for the general mathematical link between utility and information theory investigated in this paper is a simple financial market model with two kinds of small traders: less informed traders and insiders whose extra information is represented by an enlargement of the other agents' filtration. The expected logarithmic utility increment, i.e. the difference of the insider's and the less informed trader's expected logarithmic utility is described in terms of the information drift, i.e. the drift one has to eliminate in order to perceive the price dynamics as a martingale from the insider's perspective. On the one hand, we describe the information drift in a very general setting by natural quantities expressing the probabilistic better informed view of the world. This on the other hand allows us to identify the additional utility by entropy related quantities known from information theory. In particular in a complete market in which the insider has some fixed additional information during the entire trading interval, its utility increment can be represented by the Shannon information of his extra knowledge. For general markets, and in some particular examples, we provide estimates of maximal utility by information inequalities.

2000 AMS subject classifications: primary 60H30, 94A17 ; secondary 91B16, 60G44.

Key words and phrases: enlargement of filtration; logarithmic utility; utility maximization; heterogeneous information; insider model; Shannon information; information difference; entropy; differential entropy.

Introduction

A simple mathematical model of two agents on a financial markets taking their portfolio decisions on the basis of different information horizons has attracted much attention in recent years. Both agents are small, and unable to influence the price dynamics of the risky assets constituting the market. One agent just acts on the basis of the evolution of the market, the other one, the insider, possesses some additional knowledge at every instant of the continuous trading interval. This basic fact is modelled by associating two different filtrations with each agent, from which they take their portfolio decisions: the less informed agent, at time t , just has the σ -field \mathcal{F}_t , corresponding to the natural evolution of the market up to this time, at his disposal for deciding about future investments, while the insider is able to make cleverer decisions, taking his knowledge from a bigger σ -field $\mathcal{G}_t \supset \mathcal{F}_t$. We give a short selection of some among many more papers dealing with this model, just indicating the most important mathematical techniques used for its investigation. Methods are focused on martingale and stochastic control theory, and techniques of enlargement of filtrations (see Yor , Jeulin , Jacod in [JY85]), starting with the conceptual paper by Duffie, Huang [DH86], mostly in the *initial enlargement* setting, i.e. the insider gets some fixed extra information at the beginning of the trading interval. The model is successively studied on stochastic bases with increasing complexity: e.g. Karatzas, Pikovsky [PK96] on Wiener space, Gorud, Pontier [GP98] allow Poissonian noise, Biagini and Oksendal [BO03] employ anticipative calculus techniques. In the same setting, Amendinger, Becherer and Schweizer [ABS03] calculate the value of insider information from the perspective of specific utilities. Baudoin [Bau01] introduces the concept of weak additional information consisting in the knowledge of the law of some random element. Campi [Cam03] considers hedging techniques for insiders in the incomplete market setting. It is clear that the expected utility the insider is able to gain from final wealth in this simple model will be bigger than the uninformed traders' utility, for every utility function. And in fact many of the quoted papers deal with the calculation of a better informed agent's additional utility.

In Amendinger et al. [AIS98], in the setting of initial enlargements, a crucial and natural link between the additional expected logarithmic utility and information theoretic concepts was made. The insider's utility advantage is identified with the Shannon entropy of the additional information. In the same setting, Gasbarra, Valkeila [GV03] extended this link by interpreting the logarithmic utility increment by the Kullback-Leibler information of the insider's additional knowledge from the perspective of Bayesian modelling. In the environment of this utility-information paradigm the papers [Imk96], [IPW01], [Imk02], [Imk03], Corcuera et al. [CIKHN03], and Ankirchner et al. [AI04] describe additional utility, treat arbitrage questions and their interpretation in information theoretic terms in increasingly complex models of the same base structure, including some simple examples of progressive enlargements. It is clear that utility concepts different from the logarithmic one correspond on the information theoretic side to the generalized entropy concepts of f -divergences.

In this paper we shall continue the investigation of mathematical questions related to the link between utility and information theory in the most general setting of enlargements of filtrations: besides assuming eventually that the base space be standard, to ensure the existence of regular conditional probabilities, we shall let the filtration of the better informed

agent to just contain the one of the natural evolution of knowledge. To concentrate on just one kind of entropy in this general setting, we shall consider logarithmic utility throughout. In this framework, Ankirchner et al. [AI04], calculates the optimal utility of an agent from the intrinsic point of view of his (general) filtration, and relates finiteness of expected utility via the (NFLVR) condition to the characterization of semimartingales by the theorem of Dellacherie-Meyer-Mokobodski. The compensator in the Doob-Meyer decomposition of underlying asset price processes with respect to the agent's filtration is dominated by the *information drift process*. In this paper we shall give a general analysis of the nature of this process, and in particular relate it to measuring the difference of the information residing in the two filtrations of the basic agents, which of course has to be independent of the particular price dynamics. The basic and key observation we start with in section 2 identifies the information drift process with Radon-Nikodym densities of the stochastic kernel in an integral representation of the conditional probability process related to the small filtration and the conditional probability process itself, both restricted to the σ -fields of the bigger filtration. In this perspective, the information drift is just the drift to subtract from the price process from the point of view of the better informed agent to view the price as a local martingale after changing the measure according to Girsanov. This observation yields in addition the access to an identification of the additional utility described by an "energy integral" of the information drift by the information difference of the two filtrations in terms of Shannon entropy notions in section 5, which is independent of concrete price dynamics of the financial market.

In more details, the paper is organized as follows. In the preparatory section 1, we recall the main results about the connection between finite utility filtrations, properties of the price dynamics from the perspective of different agents, and properties of the information drift from Ankirchner et al. [AI04]. In the crucial section 2 (Theorems 2.6 and 2.10) properties of the conditional probability processes with respect to the agents' filtrations and the information drift process are investigated at depth, and lead to the fundamental identification of the information drift by subjective conditional probability quantities. The description of the additional utility in terms of entropy notions is more easily obtained, if the additional information in the bigger filtration comes in discrete bits along a sequence of partitions of the trading interval, leading to stepwise "initial enlargements" which ultimately converge to the big filtration as the mesh of the partitions shrinks to 0. This is done in section 5 (Theorem 5.9), after being prepared in sections 3 and 4 by a general investigation of the convergence properties of information drifts going along with the convergence of such discretized enlargements to the big filtration. In the final section 6, general facts known from Shannon information theory (see Ihara [Iha93]) are applied to estimate the expected maximal logarithmic utility of a better informed agent via the identification Theorem of section 5, in several particular cases. Entropy maximizing properties of Gaussian random variables play an important role.

1 Preliminaries

In this preparatory section we define the financial market model and recall some basic facts about maximal utility hedging. Our favorite utility function will be the logarithmic one, for which we will then compare the maximal expected utilities of agents on the market who act on the background of asymmetric information. Recalling a result from [AI04], we will describe the utility increment of a better informed agent by the respective *information drift* of the agents' filtrations.

Let (Ω, \mathcal{F}, P) be a probability space with a filtration $(\mathcal{F}_t)_{0 \leq t \leq T}$, where $T > 0$ is a fixed time horizon. We consider a financial market with one non-risky asset of interest rate normalized to 0, and one risky asset with price S_t at time $t \in [0, T]$. We assume that S is a continuous (\mathcal{F}_t) -semimartingale with values in \mathbb{R} and write \mathcal{A} for the set of all S -integrable and (\mathcal{F}_t) -predictable processes. If $\theta \in \mathcal{A}$, then we denote by $(\theta \cdot S)$ the usual stochastic integral process. For all $x > 0$ we interpret

$$x + (\theta \cdot S)_t, \quad 0 \leq t \leq T,$$

as the wealth process of a trader possessing an initial wealth x and choosing the investment strategy θ on the basis of his knowledge horizon corresponding to the filtration (\mathcal{F}_t) .

Throughout this paper we will suppose the preferences of the agents to be described by the logarithmic utility function. Furthermore we suppose that the traders' total wealth has always to be strictly positive, i.e. for all $t \in [0, T]$

$$x + (\theta \cdot S)_t > 0 \quad \text{a.s.} \tag{1}$$

Strategies θ satisfying equation (1) will be called *x-superadmissible*. The agents want to maximize their expected logarithmic utility from their wealth at time T . So we are interested in the exact value of

$$u(x) = \sup\{E \log(x + (\theta \cdot S)_T) : \theta \in \mathcal{A} \text{ } x\text{-superadmissible}\}.$$

Sometimes we will write $u_{\mathcal{F}}(x)$, in order to precise the underlying filtration. The expected logarithmic utility of the agent can be calculated easily, if one has a semimartingale decomposition of the form

$$S_t = M_t + \int_0^t \alpha_s d\langle M, M \rangle_s, \tag{2}$$

where α is a predictable process. Such a decomposition is given for a large class of semimartingales. For example if S satisfies the property (NFLVR), then it may be decomposed as in equation (2) (see [DS95]). As is shown in a forthcoming PhD thesis [Ank04], finiteness of $u(x)$ implies already such a decomposition to exist. Hence a decomposition as in (2) may be given even in cases where arbitrage exists. We state Theorem 4.4 of [AI04].

Proposition 1.1. *Suppose S can be decomposed into $S = M + \alpha \cdot \langle M, M \rangle$. Then for any $x > 0$ the following equation holds*

$$u(x) = \log x + \frac{1}{2} E \int_0^T \alpha_s^2 d\langle M, M \rangle_s. \tag{3}$$

This proposition motivates the following definition.

Definition 1.2. A filtration (\mathcal{G}_t) is called *finite utility filtration* for S , if S is a (\mathcal{G}_t) -semimartingale with decomposition $dS = dM + \alpha \cdot d\langle M, M \rangle$, where α is (\mathcal{G}_t) -predictable and belongs to $L^2(M)$, i.e. $E \int_0^T \alpha^2 d\langle M, M \rangle < \infty$. We write

$$\mathbb{F} = \{(\mathcal{H}_t) \supset (\mathcal{F}_t) \mid (\mathcal{H}_t) \text{ is a finite utility filtration for } S\}.$$

We now compare two traders who take their portfolio decisions not on the basis of the same filtration, but on the basis of different information flows represented by the filtrations (\mathcal{G}_t) and (\mathcal{H}_t) respectively. Suppose that both filtrations (\mathcal{G}_t) and (\mathcal{H}_t) are finite utility filtrations. We denote by

$$S = M + \alpha \cdot \langle M, M \rangle \tag{4}$$

the semimartingale decomposition with respect to (\mathcal{G}_t) and by

$$S = N + \beta \cdot \langle N, N \rangle \tag{5}$$

the decomposition with respect to (\mathcal{H}_t) . Obviously,

$$\langle M, M \rangle = \langle S, S \rangle = \langle N, N \rangle$$

and therefore the utility difference is equal to

$$u_{\mathcal{H}}(x) - u_{\mathcal{G}}(x) = \frac{1}{2} E \int_0^T (\beta^2 - \alpha^2) d\langle M, M \rangle.$$

Furthermore, the equations (4) and (5) imply

$$M = N - (\alpha - \beta) \cdot \langle M, M \rangle \quad a.s. \tag{6}$$

If $\mathcal{G}_t \subset \mathcal{H}_t$ for all $t \geq 0$, equation (6) can be interpreted as the semimartingale decomposition of M with respect to (\mathcal{H}_t) . In this case one can show that the utility difference depends only on the process $\mu = \alpha - \beta$. We therefore use the following notion.

Definition 1.3. Let (\mathcal{G}_t) be a finite utility filtration and $S = M + \alpha \cdot \langle M, M \rangle$ the Doob-Meyer decomposition of S with respect to (\mathcal{G}_t) . Suppose that (\mathcal{H}_t) is a filtration such that $\mathcal{G}_t \subset \mathcal{H}_t$ for all $t \in [0, T]$. The (\mathcal{H}_t) -adapted measurable process μ satisfying

$$M - \int_0^\cdot \mu_t d\langle M, M \rangle_t \quad \text{is a } (\mathcal{H}_t) \text{ - local martingale}$$

is called *information drift* (see [Imk03]) of (\mathcal{H}_t) with respect to (\mathcal{G}_t) .

The following Proposition relates the information drift to the expected logarithmic utility increment.

Proposition 1.4. *Let (\mathcal{G}_t) and (\mathcal{H}_t) be two finite utility filtrations such that $\mathcal{G}_t \subset \mathcal{H}_t$ for all $t \in [0, T]$. If μ is the information drift of (\mathcal{H}_t) w.r.t. (\mathcal{G}_t) , then we have*

$$u_{\mathcal{H}}(x) - u_{\mathcal{G}}(x) = \frac{1}{2} E \int_0^T \mu^2 d\langle M, M \rangle.$$

Proof. See Theorem 5.4 in [AI04]. □

So far we only required the information drift to be measurable and adapted. Due to the continuity of S we have the following.

Proposition 1.5. *The information drift, provided it exists, may be chosen to be predictable.*

Proof. Suppose μ is a measurable and (\mathcal{G}_t) -adapted process such that

$$M - \int_0^\cdot \mu_t d\langle M, M \rangle_t$$

is a (\mathcal{G}_t) -local martingale. We denote by ${}^p\mu$ the predictable projection of μ with respect to (\mathcal{G}_t) . We will show that $M - {}^p\mu \cdot \langle M, M \rangle$ remains a (\mathcal{G}_t) -local martingale.

Let τ be stopping time localizing M such that M^τ , the martingale M stopped at τ , is bounded. To simplify notation we assume $M^\tau = M$. Let $0 \leq s < t$, $A \in \mathcal{G}_s$ and $\varepsilon > 0$. Then

$$\begin{aligned} E(1_A(M_t - M_{s+\varepsilon})) &= E\left(1_A \int_{s+\varepsilon}^t \mu_r d\langle M, M \rangle_r\right) \\ &= E\left(1_A E\left[\int_{s+\varepsilon}^t \mu_r d\langle M, M \rangle_r \middle| \mathcal{G}_s\right]\right) \\ &= E\left(1_A E\left[\int_{s+\varepsilon}^t {}^p\mu_r d\langle M, M \rangle_r \middle| \mathcal{G}_s\right]\right) \\ &= E\left(1_A \int_{s+\varepsilon}^t {}^p\mu_r d\langle M, M \rangle_r\right) \end{aligned}$$

(see theorem 57, chapter VI in [DM78]). By dominated convergence the left hand side of this equation converges to $E(1_A(M_t - M_s))$ as $\varepsilon \downarrow 0$. The right hand side converges by similar arguments. Hence we obtain

$$E(1_A(M_t - M_s)) = E\left(1_A \int_s^t {}^p\mu_r d\langle M, M \rangle_r\right),$$

which means that $M - {}^p\mu \cdot \langle M, M \rangle$ is a (\mathcal{G}_t) -martingale. □

We close this section by recalling some basic properties of information drifts.

Lemma 1.6. *Suppose the filtration (\mathcal{F}_t) is a finite utility filtration with respect to which the Doob-Meyer decomposition of S is given by $S = M + \alpha \cdot \langle M, M \rangle$. Let (\mathcal{H}_t) be a filtration satisfying $\mathcal{F}_t \subset \mathcal{H}_t$ for all $t \in [0, T]$ and suppose that (\mathcal{H}_t) has an information drift μ with respect to (\mathcal{F}_t) . Then the following properties hold true.*

- i) *If μ belongs to $L^2(M)$, then the maximal expected utility $u_{\mathcal{H}}(x)$ is finite for all $x > 0$.*
- ii) *\mathbb{F} is equal to the set of all filtrations containing (\mathcal{F}_t) and possessing an information drift μ with respect to (\mathcal{F}_t) such that $\mu \in L^2(M)$.*
- iii) *If (\mathcal{G}_t) is a filtration such that $\mathcal{F}_t \subset \mathcal{G}_t \subset \mathcal{H}_t$ for all $t \in [0, T]$, then there is also an information drift κ of (\mathcal{G}_t) with respect to (\mathcal{F}_t) . More precisely, κ is equal to the $L^2(M)$ -projection of μ onto the subspace of (\mathcal{G}_t) -predictable processes.*

iv) μ is orthogonal to α , i.e. we have $E \int_0^T \mu_s \alpha_s \langle M, M \rangle_s = 0$.

Proof. Properties i) and ii) are obvious. A proof of property iii) and iv) can be found in [AI04]. \square

2 General enlargements

Assume again that the price process S is a semimartingale of the form

$$S = M + \alpha \cdot \langle M, M \rangle$$

with respect to some (finite utility) filtration (\mathcal{G}_t) . And for simplicity of notation suppose in this section that time horizon is infinite, i.e. $T = \infty$. We shall aim at describing the relative information drift α by basic quantities related to the conditional probabilities of the larger σ -algebras \mathcal{G}_t with respect to the smaller ones $\mathcal{F}_t, t \geq 0$. Roughly, modulo some tedious technical details to be specified below, the relationship is as follows. Suppose for all $t \geq 0$ there is a regular conditional probability $P_t(\cdot, \cdot)$ of \mathcal{F} given \mathcal{F}_t , which can be decomposed into a martingale component orthogonal to M , plus a component possessing a stochastic integral representation with respect to M with a kernel function $k_t(\cdot, \cdot)$. Then we shall see, that provided α is square integrable with respect to $d\langle M, M \rangle \otimes P$, the kernel function at t will be a signed measure in its set variable which is absolutely continuous with respect to the conditional probability, if restricted to \mathcal{G}_t , and α coincides with their Radon-Nikodym density.

We shall even be able to show that this relationship also makes sense in the reverse direction. Roughly, if absolute continuity of the stochastic integral kernel with respect to the conditional probabilities holds, and the Radon-Nikodym density is square integrable, the latter will turn out to provide an information drift α in a Doob-Meyer decomposition of S in the larger filtration.

We shall finish the section with an illustration of this fundamental relationship by discussing some simple examples of particularly enlarged filtrations.

The discussion of the details of this fundamental relationship requires some care with the complexity of the underlying filtrations and state spaces. Of course, the need to work with conditional probabilities first of all confines us to spaces on which they exist. Let therefore (Ω, \mathcal{F}, P) be a standard Borel probability space (see [Par77]) with a filtration $(\mathcal{F}_t^0)_{t \geq 0}$ consisting of countably generated σ -algebras, and M a (\mathcal{F}_t^0) -local martingale. We will also deal with the smallest right-continuous and completed filtration containing (\mathcal{F}_t^0) , which we denote by (\mathcal{F}_t) . We suppose that \mathcal{F}_0 is trivial and that every (\mathcal{F}_t) -local martingale has a continuous modification. Since \mathcal{F}_t^0 is a subfield of a standard Borel space there exist regular conditional probabilities P_t relative to the σ -algebras \mathcal{F}_t^0 . Then for any set $A \in \mathcal{F}$ the process

$$(t, \omega) \mapsto P_t(\omega, A)$$

is an (\mathcal{F}_t^0) -martingale with a continuous modification (see e.g. Theorem 4, chapter VI in [DM78]). Note that the modification may not be adapted to (\mathcal{F}_t^0) , but only to (\mathcal{F}_t) .

Furthermore it is no problem to assume that the processes $P_t(\cdot, A)$ are modified in a way such that $P_t(\omega, \cdot)$ remains a measure on \mathcal{F} for P_M -almost all (ω, t) , where P_M is a measure on $\Omega \times \mathbb{R}_+$ defined by $P_M(\Gamma) = E \int_0^\infty 1_\Gamma(\omega, t) d\langle M, M \rangle_t$, $\Gamma \in \mathcal{F} \otimes \mathcal{B}_+$.

It is known that each of these martingales may be uniquely written (see e.g. [RY99], Chapter V)

$$P_t(\cdot, A) = P(A) + \int_0^t k_s(\cdot, A) dM_s + L_t^A, \quad (7)$$

where $k(\cdot, A)$ is (\mathcal{F}_t) -predictable and L^A satisfies $\langle L^A, M \rangle = 0$.

Now let (\mathcal{G}_t^0) be another filtration on (Ω, \mathcal{F}, P) satisfying

$$\mathcal{F}_t^0 \subset \mathcal{G}_t^0$$

for all $0 \leq t \leq T$. We assume that each σ -field \mathcal{G}_t^0 is generated by a countable number of sets and we denote by (\mathcal{G}_t) the smallest right-continuous and completed filtration containing (\mathcal{G}_t^0) . It is clear that each σ -field in the left-continuous filtration (\mathcal{G}_{t-}^0) is also generated by a countable number of sets. We claim that the existence of an information drift of (\mathcal{G}_t) relative to (\mathcal{F}_t) for the process M depends on whether the following condition is satisfied or not.

Condition 2.1. $k_t(\omega, \cdot)|_{\mathcal{G}_{t-}^0}$ is a signed measure and satisfies

$$k_t(\omega, \cdot) \Big|_{\mathcal{G}_{t-}^0} \ll P_t(\omega, \cdot) \Big|_{\mathcal{G}_{t-}^0}$$

for P_M -a.a (ω, t) .

Remark 2.2. Unfortunately we have to distinguish between the filtrations (\mathcal{F}_t^0) , (\mathcal{G}_t^0) and their extensions (\mathcal{F}_t) , (\mathcal{G}_t) . The reason for this is that the regular conditional probabilities considered exist only with respect to the smaller σ -fields. On the other side, we use stochastic integration techniques which were developed only under the assumption that the underlying filtrations satisfy the usual conditions.

Let us next state some essential properties of the Radon-Nikodym density process existing according to our condition.

Lemma 2.3. *Suppose condition 2.1 to be satisfied. Then there is a $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable process γ such that for P_M -a.a. (ω, t)*

$$\gamma_t(\omega, \omega') = \frac{dk_t(\omega, \cdot)}{dP_t(\omega, \cdot)} \Big|_{\mathcal{G}_{t-}^0}.$$

Remark 2.4. Note that $\gamma_t(\omega, \cdot)$ is \mathcal{G}_{t-} -measurable. This is due to the fact that the predictable σ -algebra does not change by taking the left-continuous version of the underlying filtration.

Proof. Let $t_i^n = \frac{i}{2^n}$ for all $n \geq 0$ and $i \geq 0$. We denote by \mathbb{T} the set of all t_i^n . It is possible to choose a family of finite partitions $(\mathcal{P}^{i,n})$ such that

- for all $t \in \mathbb{T}$ we have $\mathcal{G}_{t-}^0 = \sigma(\mathcal{P}^{i,n} : i, n \geq 0 \text{ s.t. } t_i^n = t)$,

- $\mathcal{P}^{i,n} \subset \mathcal{P}^{i+1,n}$,
- if $i < j$, $n < m$ and $i2^{-n} = j2^{-m}$, then $\mathcal{P}^{i,n} \subset \mathcal{P}^{j,m}$.

We define for all $n \geq 0$

$$\gamma_t^n(\omega, \omega') = \sum_{i \geq 0} \sum_{A \in \mathcal{P}^{i,n}} 1_{]t_i^n, t_{i+1}^n]}(t) 1_A(\omega') \frac{k_t(\omega, A)}{P_t(\omega, A)}.$$

Note that $\frac{k_t(\omega, A)}{P_t(\omega, A)}$ is (\mathcal{F}_t) -predictable and $1_{]t_i^n, t_{i+1}^n]}(t) 1_A(\omega')$ is (\mathcal{G}_t) -predictable. Hence the product of both functions, defined as a function on $\Omega^2 \times \mathbb{R}_+$, is predictable with respect to $(\mathcal{F}_t \otimes \mathcal{G}_t)$. It follows that each γ^n , and thus

$$\gamma = \liminf_{n \rightarrow \infty} \gamma^n$$

is $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable.

Now fix $t \geq 0$. We claim that $k_t(\omega, \cdot) = \int \gamma_t(\omega, \omega') P_t(\omega, d\omega')$, and hence that $\gamma_t(\omega, \cdot)$ is the density of $k_t(\omega, \cdot)$ with respect to $P_t(\omega, \cdot)$, P_M -a.s. For all $n \geq 0$ let $j = j(n)$ be the integer satisfying $t_j^n < t \leq t_{j+1}^n$ and denote by \mathcal{Q}^n the corresponding partition $\mathcal{P}^{j,n}$. Observe that (\mathcal{Q}^n) is an increasing sequence of partitions satisfying

$$\sigma(\mathcal{Q}^n : n \geq 0) = \mathcal{G}_{t-}^0$$

and hence

$$\begin{aligned} \gamma_t(\omega, \omega') &= \liminf_n \gamma_t^n(\omega, \omega') \\ &= \liminf_n \sum_{A \in \mathcal{Q}^n} 1_A(\omega') \frac{k_t(\omega, A)}{P_t(\omega, A)} \\ &= \left. \frac{dk_t(\omega, \cdot)}{dP_t(\omega, \cdot)} \right|_{\mathcal{G}_{t-}^0}. \end{aligned}$$

□

Lemma 2.5. *If $(t, \omega, \omega') \mapsto \theta_t(\omega, \omega')$ is $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable and bounded, then*

$$\int \int \int \theta_t(\omega, \omega') P_t(\omega, d\omega') d\langle M, M \rangle_t dP(\omega) = \int \int \theta_t(\omega, \omega) d\langle M, M \rangle_t dP(\omega).$$

Proof. Let $0 \leq r < s$, $A \in \mathcal{F}_r$, $B \in \mathcal{G}_r$ and

$$\theta_t(\omega, \omega') = 1_{]r, s]}(t) 1_A(\omega) 1_B(\omega').$$

Then

$$\begin{aligned} &\int \int \int \theta_t(\omega, \omega') P_t(\omega, d\omega') d\langle M, M \rangle_t dP(\omega) \\ &= \int \int_r^s 1_A(\omega) P_t(\omega, B) d\langle M, M \rangle_t dP(\omega) \\ &= \int \int_r^s 1_A(\omega) 1_B(\omega) d\langle M, M \rangle_t dP(\omega) \\ &= \int \int \theta_t(\omega, \omega) d\langle M, M \rangle_t dP(\omega), \end{aligned}$$

where the second equality holds due to results about optional projections (see Theorem 57, chapter VI, in [DM78]). By a monotone class argument this can be extended to all bounded and $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable processes. \square

Theorem 2.6. *Suppose condition 2.1 is satisfied and γ is as in Lemma 2.3. Then*

$$\alpha_t(\omega) = \gamma_t(\omega, \omega)$$

is the information drift of (\mathcal{G}_t) relative to (\mathcal{F}_t) .

Proof. Suppose τ to be a stopping time such that M^τ is a martingale. For $0 \leq s < t$ and $A \in \mathcal{G}_s^0$ we have to show

$$E[1_A(M_t^\tau - M_s^\tau)] = E\left[1_A \int_s^t \gamma_u(\omega, \omega) d\langle M, M \rangle_u^\tau\right].$$

For notational simplicity write $M^\tau = M$ and observe

$$\begin{aligned} E[1_A(M_t - M_s)] &= E[P_t(\cdot, A)(M_t - M_s)] \\ &= E\left[(M_t - M_s) \int_0^t k_u(\cdot, A) dM_u\right] + E[(M_t - M_s)L_t^A] \\ &= E\left[\int_s^t k_u(\cdot, A) d\langle M, M \rangle_u\right] \\ &= E\left[\int_s^t \int_A \gamma_u(\omega, \omega') dP_u(\omega, d\omega') d\langle M, M \rangle_u\right] \\ &= E\left[1_A(\omega) \int_s^t \gamma_u(\omega, \omega) d\langle M, M \rangle_u\right], \end{aligned}$$

where we used Lemma 2.5 in the last equation. \square

Corollary 2.7. *(\mathcal{G}_t) is a finite utility filtration if and only if*

$$\int \int \int \gamma_t^2(\omega, \omega') P_t(\omega, d\omega') d\langle M, M \rangle_t dP(\omega) < \infty.$$

Proof. This follows immediately from Lemma 2.5. \square

We now look at the problem from the reverse direction. Starting with the assumption that (\mathcal{G}_t) is a finite utility filtration, we show the validity of condition 2.1.

In the sequel, (\mathcal{G}_t) denotes a finite utility filtration and α its predictable information drift, i.e.

$$\tilde{M} = M - \int_0^\cdot \alpha_t d\langle M, M \rangle_t$$

is a (\mathcal{G}_t) -local martingale. To prove the main theorem we need the following lemma.

Lemma 2.8. *Let $0 \leq s < t$ and $\mathcal{P} = \{A_1, \dots, A_n\}$ be a finite partition of Ω into \mathcal{G}_s^0 -measurable sets. Then*

$$E \int_s^t \sum_{k=1}^n \left(\frac{k_u}{P_u}\right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \leq 4E \left(\int_s^t \alpha_u^2 d\langle M, M \rangle_u \right) < \infty.$$

Proof. Let $\mathcal{P} = \{A_1, \dots, A_n\}$ be a finite \mathcal{G}_s^0 -partition. An application of Ito's formula yields

$$\begin{aligned}
& \sum_{k=1}^n [1_{A_k} \log P_s(\cdot, A_k) - 1_{A_k} \log P_t(\cdot, A_k)] \\
&= \sum_{k=1}^n \left[- \int_s^t \frac{1}{P_u(\cdot, A_k)} 1_{A_k} dP_u(\cdot, A_k) \right. \\
&\quad \left. + \frac{1}{2} \int_s^t \frac{1}{P_u(\cdot, A_k)^2} 1_{A_k} d\langle P(\cdot, A_k), P(\cdot, A_k) \rangle_u \right] \\
&= \sum_{k=1}^n \left[- \int_s^t \frac{k_u}{P_u}(\cdot, A_k) 1_{A_k} d\tilde{M}_u - \int_s^t \frac{k_u}{P_u}(\cdot, A_k) 1_{A_k} \alpha_u d\langle M, M \rangle_u \right. \\
&\quad - \int_s^t \frac{1}{P_u(\cdot, A_k)} 1_{A_k} dL_u^{A_k} + \frac{1}{2} \int_s^t \left(\frac{k_u}{P_u} \right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \\
&\quad \left. + \frac{1}{2} \int_s^t \frac{1}{P_u(\cdot, A_k)^2} 1_{A_k} d\langle L^{A_k}, L^{A_k} \rangle_u \right] \tag{8}
\end{aligned}$$

Note that $P_t(\cdot, A_k) \log P_t(\cdot, A_k)$ is a submartingale bounded from below for all k . Hence the expectation of the left hand side in the previous equation is at most 0.

A priori it is not clear whether

$$\sum_{k=1}^n \int_s^t \frac{k_u}{P_u}(\cdot, A_k) 1_{A_k} d\tilde{M}_u$$

is integrable or not. Consider therefore for all $\varepsilon > 0$ stopping times defined by

$$\tau_k^\varepsilon = \begin{cases} \infty & \omega \notin A_k \\ \inf\{t \geq s : P_t(\cdot, A_k) \leq \varepsilon\} & \text{else} \end{cases}$$

and

$$\tau^\varepsilon = \tau_1^\varepsilon \wedge \dots \wedge \tau_n^\varepsilon.$$

Observe that $\tau^\varepsilon \rightarrow \infty$ as $\varepsilon \downarrow 0$ and that the stopped process

$$\sum_{k=1}^n \int_s^{t \wedge \tau^\varepsilon} \frac{k_u}{P_u}(\cdot, A_k) 1_{A_k} d\tilde{M}_u$$

has expectation zero, since

$$\begin{aligned}
& E \left[\left(\int_s^{t \wedge \tau^\varepsilon} \sum_{k=1}^n \frac{k_u}{P_u}(\cdot, A_k) 1_{A_k} d\tilde{M}_u \right)^2 \right] \\
&= E \left[\int_s^{t \wedge \tau^\varepsilon} \sum_{k=1}^n \left(\frac{k_u}{P_u} \right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \right] \\
&\leq \frac{1}{\varepsilon^2} E \left[\int_s^{t \wedge \tau^\varepsilon} \sum_{k=1}^n (k_u)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \right] \\
&\leq \frac{1}{\varepsilon^2} E \left[\sum_{k=1}^n \int_s^t d\langle P(\cdot, A_k), P(\cdot, A_k) \rangle_u \right] \\
&< \infty.
\end{aligned}$$

Similarly, one can show that the expectation of

$$\int_s^{t \wedge \tau^\varepsilon} \frac{1}{P_u(\cdot, A_k)} 1_{A_k} dL_u^{A_k}$$

vanishes. Consequently we may deduce from equation (8) and the Kunita-Watanabe inequality

$$\begin{aligned} & E \sum_{k=1}^n \frac{1}{2} \int_s^{t \wedge \tau^\varepsilon} \left(\frac{k_u}{P_u} \right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \\ & \leq E \sum_{k=1}^n \left[\int_s^{t \wedge \tau^\varepsilon} \frac{k_u}{P_u} (\cdot, A_k) 1_{A_k} \alpha_u d\langle M, M \rangle_u \right] \\ & \leq E \left(\int_s^{t \wedge \tau^\varepsilon} \sum_{k=1}^n \left(\frac{k_u}{P_u} \right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \right)^{\frac{1}{2}} E \left(\int_s^{t \wedge \tau^\varepsilon} \alpha_u^2 d\langle M, M \rangle_u \right)^{\frac{1}{2}}, \end{aligned}$$

which implies

$$E \int_s^{t \wedge \tau^\varepsilon} \sum_{k=1}^n \left(\frac{k_u}{P_u} \right)^2 (\cdot, A_k) 1_{A_k} d\langle M, M \rangle_u \leq 4E \left(\int_s^{t \wedge \tau^\varepsilon} \alpha_u^2 d\langle M, M \rangle_u \right).$$

Now the proof may be completed by a monotone convergence argument. \square

Let \mathbb{T} and $(\mathcal{P}^{i,n})_{i,n \geq 0}$ be a family of partitions as in the proof of Lemma 2.3. We define for all $n \geq 0$

$$Z_t^n(\omega, \omega') = \sum_{i \geq 0} \sum_{A \in \mathcal{P}^{i,n}} 1_{]t_i^n, t_{i+1}^n]}(t) 1_A(\omega') \frac{k_t(\omega, A)}{P_t(\omega, A)}.$$

Note that Z^n is $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable. We are now able to prove a converse statement to Theorem 2.6. Observe first

Lemma 2.9. *For P_M -almost all $(\omega, t) \in \Omega \times \mathbb{R}_+$ the discrete process $(Z_t^m(\omega, \cdot))_{m \geq 1}$ is an $L^2(P_t(\omega, \cdot))$ -bounded martingale.*

Proof. Every statement in the sequel is meant to hold for P_M -a.a. $(\omega, t) \in \Omega \times \mathbb{R}_+$.

Let $m \geq 0$, $l \geq 0$ and j be the natural number such that $]t_l^{m+1}, t_{l+1}^{m+1}] \subset]t_j^m, t_{j+1}^m]$. We start by proving that on $]t_l^{m+1}, t_{l+1}^{m+1}]$ we have

$$E^{P_t(\omega, \cdot)}[Z_t^{m+1}(\omega, \cdot) | \mathcal{P}^{j,m}] = Z_t^m(\omega, \cdot).$$

For this, let $B \in \mathcal{P}^{j,m}$ and $A_1, \dots, A_k \in \mathcal{P}^{l,m+1}$ such that $A_1 \cup \dots \cup A_k = B$. Note that

$$\begin{aligned} E^{P_t(\omega, \cdot)}[1_B(\cdot) Z_t^{m+1}(\omega, \cdot)] &= E^{P_t(\omega, \cdot)} \left[\sum_{i=1}^k 1_{A_i}(\cdot) \frac{k_t(\omega, A_i)}{P_t(\omega, A_i)} \right] \\ &= \sum_{i=1}^k k_t(\omega, A_i) \\ &= k_t(\omega, B) \\ &= E^{P_t(\omega, \cdot)}[1_B(\cdot) Z_t^m(\omega, \cdot)] \end{aligned}$$

on $]t_l^{m+1}, t_{l+1}^{m+1}]$. Consequently the process $(Z_t^m(\omega, \cdot))_{m \geq 1}$ is a martingale (with respect to a filtration depending on t). The martingale property implies that the sequence $\int (Z_t^n)^2(\omega, \omega') P_t(\omega, d\omega')$ is increasing, and hence, by monotone convergence,

$$\begin{aligned} & \sup_n E \int \int (Z_t^n)^2(\omega, \omega') P_u(\omega, d\omega') d\langle M, M \rangle_t \\ &= E \int \sup_n \int (Z_t^n)^2(\omega, \omega') P_u(\omega, d\omega') d\langle M, M \rangle_t. \end{aligned}$$

By Lemma 2.8 and Lemma 2.5 we have

$$\begin{aligned} & \sup_n E \int \int (Z_u^n)^2(\omega, \omega') P_u(\omega, d\omega') d\langle M, M \rangle_u \\ &= \sup_n E \int \int (Z_u^n)^2(\omega, \omega) d\langle M, M \rangle_u \\ &= \sup_n E \sum_{i \geq 0} \int_{t_i^n}^{t_{i+1}^n} \sum_{A \in \mathcal{P}^{i,n}} 1_A(\omega) \left(\frac{k_t(\omega, A)}{P_t(\omega, A)} \right)^2 d\langle M, M \rangle_u \\ &\leq 4E \left(\int \alpha_u^2 d\langle M, M \rangle_u \right) < \infty. \end{aligned}$$

This shows that $(Z^n)_{n \geq 1}$ is an $L^2(P_t(\omega, \cdot))$ -bounded martingale. \square

We now will show that k can be chosen to be a signed measure. For this we identify $P_t(\omega, \cdot)$ with another measure on a countable generator of \mathcal{G}_{t-}^0 and apply the $(\pi - \lambda)$ -Theorem.

Theorem 2.10. *The kernel k may be chosen such that*

$$\mathcal{G}_{t-}^0 \rightarrow \mathbb{R}, A \mapsto k_t(\omega, A),$$

is a signed measure which is absolutely continuous with respect to $P_t(\omega, \cdot)|_{\mathcal{G}_{t-}^0}$, for P_M -a.a. $(\omega, t) \in \Omega \times [0, \infty)$. This means that condition 2.1 is satisfied.

Proof. Lemma 2.9 implies that the martingale $(Z_t^m(\omega, \cdot))_{m \geq 1}$ is an $L^2(P_t(\omega, \cdot))$ -bounded martingale and hence, for a.a. fixed (ω, t) , $(Z_t^m(\omega, \cdot))_{m \geq 1}$ possesses a limit Z . It can be chosen to be $(\mathcal{F}_t \otimes \mathcal{G}_t)$ -predictable. Take for example

$$Z_t = \liminf_n (Z_t^n \vee 0) + \limsup_n (Z_t^n \wedge 0).$$

Now define a signed measure by

$$\tilde{k}_t(\omega, A) = \int 1_A(\omega') Z_t(\omega, \omega') dP_t(\omega, d\omega').$$

Observe that $\tilde{k}_t(\omega, \cdot)$ is absolutely continuous with respect to $P_t(\omega, \cdot)$ and that we have for all $A \in \mathcal{P}^{j,m}$ with $j2^{-m} \leq t$

$$\tilde{k}_t(\omega, A) = k_t(\omega, A)$$

for P_M -a.a. $(\omega, t) \in \Omega \times \mathbb{R}_+$. Hence

$$P_t(\omega, A) = P(A) + \int_0^t \tilde{k}_s(\omega, A) dM_s + L_t^A(\omega) \quad (9)$$

for all $A \in \bigcup_{j_2 - m \leq t} \mathcal{P}^{j,m}$. Since the LHS and both expressions on the RHS are measures coinciding on a system which is stable for intersections, equation (9) holds for all $A \in \mathcal{G}_{t-}^0$. Hence, by choosing $k_t(\cdot, A) = \tilde{k}_t(\cdot, A)$ for all $A \in \mathcal{G}_{t-}^0$, the proof is complete. \square

Remark 2.11. Since k is determined up to P_M -null sets, we may assume that $k_t(\omega, \cdot)$ is absolutely continuous relative to $P_t(\omega, \cdot)$ everywhere.

We close this section with some examples showing how (well known) information drifts can be derived explicitly based on the formalism of Theorem 2.6. To this end it is not always necessary to determine the signed measures $k_t(\omega, \cdot)$ on the whole σ -algebras \mathcal{G}_t^0 , but only on some sub- σ -fields. This is the case for example, if

$$\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \mathcal{H}_t^0, \quad 0 \leq t \leq T,$$

where (\mathcal{H}_t^0) is some countably generated filtration on (Ω, \mathcal{F}) .

Now suppose that $k_t(\omega, \cdot)$ is a signed measure on (\mathcal{H}_{t-}^0) satisfying

$$k_t(\omega, \cdot) \Big|_{\mathcal{H}_{t-}^0} \ll P_t(\omega, \cdot) \Big|_{\mathcal{H}_{t-}^0}$$

for P_M -a.a. (ω, t) . Then we can show with the arguments of the proof of Lemma 2.3 that there is an $(\mathcal{F}_t \otimes \mathcal{H}_t)$ -predictable process β such that P_M -a.e.

$$\beta_t(\omega, \omega') = \frac{dk_t(\omega, \cdot)}{dP_t(\omega, \cdot)} \Big|_{\mathcal{H}_{t-}^0}.$$

The information drift of (\mathcal{G}_t) relative to (\mathcal{F}_t) is already determined by the trace of (β_t) . For the corresponding analogue of Theorem 2.6 we shall give a more explicit statement.

Theorem 2.12. *The process*

$$\alpha_t(\omega) = \beta_t(\omega, \omega)$$

is the information drift of (\mathcal{G}_t) relative to (\mathcal{F}_t) .

Proof. Suppose T to be a stopping time such that M^T is a martingale. For $0 \leq s < t$, $A \in \mathcal{H}_s^0$ and $B \in \mathcal{F}_s^0$ we have to show

$$E [1_A 1_B (M_t^T - M_s^T)] = E \left[1_A 1_B \int_s^t \beta_u(\omega, \omega) d\langle M, M \rangle_u^T \right].$$

For simplicity assume $M^T = M$ and observe

$$\begin{aligned} E [1_A 1_B (M_t - M_s)] &= E [1_B P_t(\cdot, A) (M_t - M_s)] \\ &= E \left[1_B (M_t - M_s) \int_0^t k_u(\cdot, A) dM_u \right] + E [1_B (M_t - M_s) L_t^A] \\ &= E \left[\int_s^t 1_B k_u(\cdot, A) d\langle M, M \rangle_u \right] \\ &= E \left[\int_s^t \int_A 1_B(\omega) \beta_u(\omega, \omega') dP_u(\omega, d\omega') d\langle M, M \rangle_u \right] \\ &= E \left[1_A(\omega) 1_B(\omega) \int_s^t \beta_u(\omega, \omega) d\langle M, M \rangle_u \right], \end{aligned}$$

where we used Lemma 2.5 in the last equation. \square

Example 2.13. Let (W_t) be the standard Wiener process and (\mathcal{F}_t^0) the filtration generated by (W_t) . Moreover, let (Y_t) be a Gaussian process independent of \mathcal{F}_1 such that for each pair s, t with $0 \leq s < t$ the difference $Y_t - Y_s$ is independent of Y_t . We denote by w_t the variance of Y_t .

We enlarge our filtration by

$$\mathcal{H}_t^0 = \sigma(W_1 + Y_s : 0 \leq s \leq t) = \sigma(W_1 + Y_t) \vee \sigma(Y_t - Y_s : 0 \leq s \leq t),$$

and put $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \mathcal{H}_t^0$, $0 \leq t \leq 1$. Now observe that for all $C \in \sigma(Y_t - Y_s : 0 \leq s \leq t)$ and Borel sets $B \in \mathcal{B}(\mathbb{R})$ we have

$$\begin{aligned} P_t(\cdot, \{W_1 + Y_t \in B\} \cap C) &= P(C) \int 1_B(x + W_1 - W_t + Y_t) dP \Big|_{x=W_t} \\ &= P(C) \int 1_B(y + x) \phi_{1-t+w_t}(y) dy \Big|_{x=W_t} \\ &= P(C) \int_B \phi_{1-t+w_t}(y - W_t) dy, \quad 0 \leq t < 1, \end{aligned}$$

where

$$\phi_v(y) = \frac{1}{(2\pi v)^{\frac{1}{2}}} e^{-\frac{y^2}{2v}}.$$

Now observe that $f(x, t) = P(C) \int_B \phi_{1-t+w_t}(y - x) dy$ is differentiable in x and satisfies

$$\frac{\partial}{\partial x} f(x, t) = P(C) \int_B \frac{y - x}{1 - t + w_t} \phi_{1-t+w_t}(y - x) dy$$

for all $0 \leq t < 1$ and $x \in \mathbb{R}$. By Ito's formula

$$P_t(\cdot, \{W_1 + Y_t \in B\} \cap C) = f(0, 0) + \int_0^t \frac{\partial}{\partial x} f(W_s, s) dW_s + A_t, \quad 0 \leq t < 1,$$

where A is a process of bounded variation. Note that A is also a martingale, and thus $A = 0$. Hence

$$\begin{aligned} &k_t(\cdot, \{W_1 + Y_t \in B\} \cap C) \\ &= P(C) \int_B \frac{y - W_t}{1 - t + w_t} \phi_{1-t+w_t}(y - W_t) dy \\ &= P(C) \int 1_B(y + x) \frac{y + x - x}{1 - t + w_t} \phi_{1-t+w_t}(y) dy \Big|_{x=W_t(\omega)} \\ &= \int_{\{W_1 + Y_t \in B\} \cap C} \frac{W_1(\omega') + Y_t(\omega') - W_t(\omega)}{1 - t + w_t} dP_t(\omega, d\omega') \end{aligned}$$

As a consequence

$$\beta_t(\omega, \omega') = \frac{k_t(\omega, d\omega')}{P_t(\omega, d\omega')} \Big|_{\mathcal{H}_t^0} = \frac{W_1(\omega') + Y_t(\omega') - W_t(\omega)}{1 - t + w_t},$$

and by Theorem 2.12,

$$W_t - \int_0^t \frac{W_1 + Y_s - W_s}{1 - s + w_s}, \quad 0 \leq t < 1,$$

is a martingale relative to (\mathcal{G}_t) .

Similar examples can be found in [CIKHN03], where the information drifts are derived in a completely different way, though.

Example 2.14. Let (W_t) be the standard Wiener process and (\mathcal{F}_t) the Wiener filtration. We use the abbreviation $W_t^* = \sup_{0 \leq s \leq t} W_s$ and consider the filtration enlarged by the random variable $G = 1_{[0,c]}(W_1^*)$, $c > 0$. Again we want to apply Theorem 2.12 in order to obtain the information drift of $\mathcal{G}_t = \mathcal{F}_t \vee \sigma(G)$. To this end let $Z_t = \sup_{t \leq r \leq 1} (W_r - W_t)$ and denote by p_t the density of Z_t , $0 \leq t < 1$. Now,

$$\begin{aligned} P_t(\cdot, G = 1) &= P(W_t^* \vee W_t + Z_t \leq c | \mathcal{F}_t) \\ &= \int 1_{[0,c]}(y \vee x + Z_t) dP \Big|_{x=W_t, y=W_t^*} \\ &= 1_{[0,c]}(y) \int_0^{c-x} p_t(z) dz \Big|_{x=W_t, y=W_t^*}, \end{aligned}$$

for all $0 \leq t < 1$. Note that $F(x, y, t) = 1_{[0,c]}(y) \int_0^{c-x} p_t(z) dz$ is differentiable in x for all $0 \leq t < 1$ and $x \in \mathbb{R}$, and by Ito's formula

$$P_t(\cdot, G = 1) = F(0, 0, 0) + \int_0^t \frac{\partial}{\partial x} F(W_s, W_s^*, s) dW_s + A_t, \quad 0 \leq t < 1,$$

where A is a process of bounded variation. Hence

$$k_t(\cdot, G = 1) = \frac{\partial}{\partial x} F(W_t, W_t^*, t), \quad 0 \leq t < 1.$$

Similarly, we have

$$P_t(\cdot, G = 0) = H(W_t, W_t^*, t), \quad 0 \leq t < 1,$$

and

$$k_t(\cdot, G = 0) = \frac{\partial}{\partial x} H(W_t, W_t^*, t), \quad 0 \leq t < 1,$$

where

$$H(x, y, t) = 1_{(c,\infty)}(y) + 1_{[0,c]}(y) \int_{c-x}^{\infty} p_t(z) dz.$$

As a consequence

$$\begin{aligned} \beta_t(\omega, \omega') &= \frac{k_t(\omega, d\omega')}{P_t(\omega, d\omega')} \Big|_{\sigma(G)} \\ &= 1_{\{1\}}(G(\omega')) \frac{\partial}{\partial x} \log F(W_t(\omega), W_t^*(\omega'), t) \\ &\quad + 1_{\{0\}}(G(\omega')) \frac{\partial}{\partial x} \log H(W_t(\omega), W_t^*(\omega'), t), \quad 0 \leq t < 1. \end{aligned}$$

3 Monotone convergence of information drifts

In the preceding section we established a general relationship between the information drift and the regular conditional probabilities of filtrations. In this framework the knowledge of

the better informed agent is described by a general enlarged filtration (\mathcal{G}_t) of (\mathcal{F}_t) . We shall now consider the question whether this situation may be well approximated by "stepwise initial" enlargements, for which we take $\mathcal{F}_t \vee \mathcal{G}_{t_i-}$ for $t \in [t_i, t_{i+1})$, if the family $(t_i)_{0 \leq i \leq n}$ is a partition of \mathbb{R}_+ . One particularly important question in this context concerns the behavior of the information drifts along such a sequence of discretized enlargements. Of course we expect some convergence of the drifts. We shall establish this fact rigorously in the following section. In the present section, we shall prepare the treatment of this problem by solving a somewhat more general problem. Let $(\mathcal{G}_t^n)_{n \in \mathbb{N}}$ be an increasing sequence of finite utility filtrations and $\sup_n u_{\mathcal{G}_t^n}(x)$ be finite. We will show that then the smallest filtration containing every (\mathcal{G}_t^n) is also a finite utility filtration.

Since we will not deal with regular conditional probabilities in this section, it is not necessary to require our probability space (Ω, \mathcal{F}) to be standard.

We use the terminology of Revuz and Yor [RY99]: $H^2(\mathcal{F}_t)$ denotes the set of L^2 -bounded continuous (\mathcal{F}_t) -martingales, i.e. the space of continuous (\mathcal{F}_t, P) -martingales M such that

$$\sup_{t \geq 0} E(M_t^2) < \infty.$$

We need the following characterization of $H^2(\mathcal{F}_t)$.

Lemma 3.1. *(Proposition 1.23 in [RY99]) A continuous (\mathcal{F}_t) -local martingale belongs to $H^2(\mathcal{F}_t)$ if and only if the following two conditions hold*

- i) $E(M_0^2) < \infty$,
- ii) $E(\langle M, M \rangle_\infty) < \infty$.

The properties i) and ii) are independent of the filtration considered. This is due to the fact that the quadratic variation of M does not change under a new filtration (\mathcal{G}_t) for which M is still a semimartingale. We therefore have

Lemma 3.2. *Suppose $M \in H^2(\mathcal{F}_t)$. Let (\mathcal{G}_t) be a filtration such that M is still a (\mathcal{G}_t) -semimartingale. If*

$$M = \tilde{M} + A$$

is a Doob-Meyer decomposition with respect to (\mathcal{G}_t) with $A_0 = 0$, then \tilde{M} belongs to $H^2(\mathcal{G}_t)$.

Proof. Notice that $\tilde{M}_0 = M_0$ and $\langle M, M \rangle = \langle \tilde{M}, \tilde{M} \rangle$. The claim follows now by applying lemma 3.1 twice. \square

Now let M be a continuous (\mathcal{F}_t) -local martingale and $(\mathcal{G}_t^n)_{n \geq 1}$ an increasing sequence of filtrations, i.e. for all $t \geq 0$ we have

$$\mathcal{F}_t \subset \mathcal{G}_t^1 \subset \dots \subset \mathcal{G}_t^n \subset \mathcal{G}_t^{n+1} \subset \dots$$

We assume that for all $n \geq 1$ the process M is a (\mathcal{G}_t^n) -semimartingale with Doob-Meyer decomposition of the form

$$M = M^n + \int_0^\cdot \mu_s^n d\langle M, M \rangle_s,$$

where μ^n is (\mathcal{G}_t^n) -predictable. We then have the following asymptotic property.

Lemma 3.3. *If the processes $(\mu^n)_{n \in \mathbb{N}}$ converge to some μ in $L^2(M)$, then*

$$M - \int_0^\cdot \mu_s d\langle M, M \rangle_s$$

is a local martingale with respect to $\mathcal{G}_t = \bigvee_{n \geq 1} \mathcal{G}_t^n, t \geq 0$.

Proof. Suppose the stopping time τ reduces M such that M^τ is a bounded martingale. Note that lemma 3.2 implies that the stopped processes $(M^n)^\tau$ are (\mathcal{G}_t) -martingales.

For simplicity we assume $M^\tau = M$. Choose a constant $C > 0$ such that

$$|M| \leq C,$$

and

$$E \int_0^\infty (\mu_s^n)^2 d\langle M, M \rangle_s \leq C^2$$

for all $n \geq 1$. Now let $\varepsilon > 0, 0 \leq s < t$ and $A \in \mathcal{G}_s$. It suffices to show

$$\left| E [1_A(M_t - M_s)] - E \left[1_A \int_s^t \mu_s d\langle M, M \rangle_s \right] \right| \leq \varepsilon.$$

We start by choosing n_0 such that

$$\|\mu^n - \mu\|_{L^2(M)} \leq \frac{\varepsilon}{4\sqrt{E(\langle M, M \rangle_\infty)}}$$

for all $n \geq n_0$.

Note that $\bigcup_{n \geq n_0} \mathcal{G}_s^n$ is an algebra generating the σ -algebra $\bigvee_{n \geq 1} \mathcal{G}_s^n = \mathcal{G}_s = \bigvee_{n \geq n_0} \mathcal{G}_s^n$. Hence we can find a sequence $(A_i)_{i \in \mathbb{N}}$ of sets in $\bigcup_{n \geq n_0} \mathcal{G}_s^n$ such that $P(A \Delta A_i) \rightarrow 0$. A subsequence of $(1_{A_i})_{i \in \mathbb{N}}$ converges to 1_A almost surely and therefore we may choose $n \geq n_0$ and $\tilde{A} \in \mathcal{G}_s^n$ satisfying $P(\tilde{A} \Delta A) \leq (\frac{\varepsilon}{4C})^2$ and

$$\left(E \int_s^t (1_A - 1_{\tilde{A}})^2 \langle M, M \rangle \right)^{\frac{1}{2}} \leq \frac{\varepsilon}{4C}.$$

Hence we have

$$\begin{aligned} |E [1_A(M_t - M_s)] - E [1_{\tilde{A}}(M_t - M_s)]| &\leq |E [(1_{\tilde{A}} - 1_A)(M_t - M_s)]| \\ &\leq P(\tilde{A} \Delta A)^{\frac{1}{2}} (E(M_t - M_s)^2)^{\frac{1}{2}} \\ &\leq \frac{\varepsilon}{2}. \end{aligned}$$

By applying the Kunita-Watanabe inequality we get for $n \geq n_0$

$$\begin{aligned}
& \left| E \left[1_A \int_s^t \mu_u d\langle M, M \rangle_u \right] - E \left[1_{\bar{A}} \int_s^t \mu_u^n d\langle M, M \rangle_u \right] \right| \\
& \leq \left| E \left[1_A \int_s^t (\mu_u - \mu_u^n) d\langle M, M \rangle_u \right] \right| + \left| E \left[(1_A - 1_{\bar{A}}) \int_s^t \mu_u^n d\langle M, M \rangle_u \right] \right| \\
& \leq \left(E \int_s^t 1_A d\langle M, M \rangle \right)^{\frac{1}{2}} \left(E \int_0^\infty (\mu_s - \mu_s^n)^2 d\langle M, M \rangle_s \right)^{\frac{1}{2}} \\
& \quad + \left(E \int_s^t (1_A - 1_{\bar{A}})^2 d\langle M, M \rangle \right)^{\frac{1}{2}} \left(E \int_0^\infty (\mu_s^n)^2 d\langle M, M \rangle_s \right)^{\frac{1}{2}} \\
& \leq (E\langle M, M \rangle)^{\frac{1}{2}} \|\mu - \mu^n\|_{L^2(M)} + \frac{\varepsilon}{4} \\
& \leq \frac{\varepsilon}{2},
\end{aligned}$$

and thus

$$\begin{aligned}
& \left| E [1_A(M_t - M_s)] - E \left[1_A \int_s^t \mu_u d\langle M, M \rangle_u \right] \right| \\
& \leq \left| E [1_A(M_t - M_s)] - E [1_{\bar{A}}(M_t - M_s)] \right| \\
& \quad + \left| E [1_{\bar{A}}(M_t - M_s)] - E \left[1_{\bar{A}} \int_s^t \mu_u^n d\langle M, M \rangle_u \right] \right| \\
& \quad + \left| E \left[1_{\bar{A}} \int_s^t \mu_u^n d\langle M, M \rangle_u \right] - E \left[1_A \int_s^t \mu_u d\langle M, M \rangle_u \right] \right| \\
& \leq \frac{\varepsilon}{2} + 0 + \frac{\varepsilon}{2} = \varepsilon.
\end{aligned}$$

□

In the remainder of this section we give some sufficient criteria for a sequence of information drifts to converge in $L^2(M)$. We start by the following observation.

Lemma 3.4. *For $n, m \geq 1$ with $m \geq n \geq 1$ the processes μ^n and $\mu^m - \mu^n$ are orthogonal in the Hilbert space $L^2(M)$, i.e.*

$$\mathbb{E} \left[\int_0^\infty \mu^n (\mu^m - \mu^n) d\langle M, M \rangle \right] = 0.$$

Proof. Let $1 \leq n \leq m$. Recall that for all $i \geq 1$

$$\tilde{M}^i = M - \int_0^\cdot \mu_t^i d\langle M, M \rangle_t \tag{10}$$

is a local \mathcal{G}_t^i -martingale. Then

$$\int_0^\infty \mu^n (\mu^m - \mu^n) d\langle M, M \rangle = (\mu^n \cdot \tilde{M}^m)_\infty - (\mu^n \cdot \tilde{M}^n)_\infty.$$

Note that μ^n is (\mathcal{G}_t^n) and (\mathcal{G}_t^m) -predictable. Moreover, $(\mu^n \cdot \tilde{M}^n)_t$ and $(\mu^n \cdot \tilde{M}^m)_t$ are L^2 -martingales w.r.t. to (\mathcal{G}_t^n) and (\mathcal{G}_t^m) respectively. Therefore,

$$\mathbb{E} \left[\int_0^\infty \mu^n (\mu^m - \mu^n) d\langle M, M \rangle \right] = 0.$$

□

We are now in a position to prove the main result of the section.

Theorem 3.5. *If $\sup_{n \geq 1} \|\mu^n\|_{L^2(M)}^2 < \infty$, then (μ^n) converges in $L^2(M)$ to a process μ . Moreover,*

$$M - \int_0^\cdot \mu d\langle M, M \rangle$$

is a local martingale with respect to $\mathcal{G}_t = \bigvee_{n \geq 1} \mathcal{G}_t^n, t \geq 0$.

Proof. Set $c = \sup_{n \geq 1} \|\mu^n\|_{L^2(M)}^2$. By the previous lemma for $m \geq n \geq 1$

$$\|\mu^m\|_{L^2(M)}^2 = \|\mu^n\|_{L^2(M)}^2 + \|\mu^m - \mu^n\|_{L^2(M)}^2.$$

Thus $c = \lim_{n \rightarrow \infty} \|\mu^n\|_{L^2(M)}^2$ and

$$\|\mu^m - \mu^n\|_{L^2(M)}^2 = \|\mu^m\|_{L^2(M)}^2 - \|\mu^n\|_{L^2(M)}^2 \leq c - \|\mu^n\|_{L^2(M)}^2 \rightarrow 0$$

as $n \rightarrow \infty$. Therefore $\{\mu^n\}_{n \geq 1}$ is a Cauchy sequence in $L^2(M)$. By completeness of $L^2(M)$, there exists a unique (\mathcal{G}_t) -predictable process $\mu^0 \in L^2(M)$ such that $\lim_{n \rightarrow \infty} \mu^n = \mu^0$ in $L^2(M)$. By lemma 3.3 the process $M - \int \mu^0 d\langle M, M \rangle$ is a (\mathcal{G}_t) -local martingale. □

4 Continuous and initial enlargements

In this section we relate general enlargements of filtrations to "initial enlargements" along discrete partitions of $[0, T]$, for finite horizon T . The knowledge of the insider is modeled by an arbitrary filtration $(\mathcal{G}_t)_{t \in [0, T]}$, satisfying $\mathcal{G}_t \supset \mathcal{F}_t, 0 \leq t \leq T$. For $s \in [0, T]$ we set

$$\mathcal{G}_t^s = \begin{cases} \mathcal{F}_t, & t < s \\ \mathcal{F}_t \vee \mathcal{G}_{s-}, & t \geq s. \end{cases}$$

Again the analysis of this section does not require our probability space (Ω, \mathcal{F}) to be standard.

Remark 4.1. In the case where the σ -field $\mathcal{G}_{s-}, s \in [0, T]$, is generated by a countable number of events, say $(A_n)_{n \in \mathbb{N}}$, the enlarged filtration \mathcal{G}_t^s can be viewed as initial enlargement at time s in the classical sense. In that case $\mathcal{G}_{s-} = \sigma(1_{A_n} : n \in \mathbb{N})$ and one has for $t \in [0, T]$,

$$\mathcal{F}_t \vee \mathcal{G}_{s-} = \mathcal{F}_t \vee \sigma(1_{A_n} : n \in \mathbb{N}).$$

The set $\{0, 1\}^{\mathbb{N}}$ can be endowed with a metric so that it becomes a Polish space with corresponding Borel- σ -field $\mathcal{B}(\{0, 1\})^{\otimes \mathbb{N}}$. Hence, the filtration (\mathcal{G}_t^s) can be seen as initial enlargement at time s induced by the random variable $G : \Omega \rightarrow \{0, 1\}^{\mathbb{N}}, \omega \mapsto (1_{A_n}(\omega))_{n \in \mathbb{N}}$. In particular, the standard theory of initial enlargements is applicable.

In the following, we assume that (\mathcal{G}_t^s) is for arbitrary $s \in [0, T]$ a finite utility filtration. Let for $0 \leq s \leq t \leq T$

$$\pi_0([0, s] \times (t, T]) := F(s, t) := \frac{1}{2} E \int_t^T (\mu_r^s)^2 d\langle M, M \rangle_r,$$

where μ^s is a (\mathcal{G}_t^s) -information drift. π_0 is defined on the semiring $J := \{[0, s] \times (t, T] : s \leq t\}$ over $D = \{(s, t) \in \mathbb{R}^2 : 0 \leq s < t \leq T\}$.

Lemma 4.2. *There exists a unique measure π on the Borel sets $\mathcal{B}(D)$ of D satisfying $\pi|_J = \pi_0$.*

Proof. In order to show that π_0 can be extended to a unique measure on $\mathcal{B}(D)$ it suffices to verify the following statements (see for instance Elstrodt [Els96] Satz 4.5, Paragraph 4). For any $(s, t) \in \overline{D}$ and any sequence $(s_n, t_n)_{n \in \mathbb{N}}$ in \overline{D} with $s_n \leq s$, $t_n \geq t$ and $\lim_{n \rightarrow \infty} (s_n, t_n) = (s, t)$ we have $\lim_{n \rightarrow \infty} F(s_n, t_n) = F(s, t)$. Moreover, $F(s_n, t_n) \leq F(s, t) < \infty$.

Let s_n, t_n, s and t as above. Without loss of generality we assume that (s_n) is monotonically increasing. For $u \in [t, T]$ we consider the filtrations $(\mathcal{G}_r^{s_n})_{r \in [u, T]}$, $n \in \mathbb{N}$, over the time interval $[u, T]$. The filtrations are monotonically increasing with $\bigvee_{n \in \mathbb{N}} \mathcal{G}_r^{s_n} = \mathcal{G}_r^s$, $r \in [u, T]$. Since $(\mu_r^{s_n})_{r \in [u, T]}$ are (\mathcal{G}^{s_n}) -information drifts, it follows (in analogy to lemma 3.4) that

$$E \int_u^T (\mu^s - \mu^{s_n}) \mu^{s_n} d\langle M, M \rangle = 0.$$

In particular,

$$E \int_u^T (\mu^{s_n})^2 d\langle M, M \rangle \leq E \int_t^T (\mu^s)^2 d\langle M, M \rangle < \infty.$$

By theorem 3.5 the processes $(\mu_r^{s_n})_{r \in [u, T]}$ converge to the information drift $(\mu_r^s)_{r \in [u, T]}$ in $L^2(M; [u, T])$. Therefore, for any $u \in (t, T]$,

$$\liminf_{n \rightarrow \infty} E \int_{t_n}^T (\mu^{s_n})^2 d\langle M, M \rangle \geq E \int_u^T (\mu^s)^2 d\langle M, M \rangle.$$

Due to the continuity of M the right hand side of the previous equation tends to $E \int_t^T (\mu^s)^2 d\langle M, M \rangle$ as $u \downarrow t$. Consequently, we obtain $\lim_{n \rightarrow \infty} F(s_n, t_n) = F(s, t)$. \square

The measure π describes the utility increase by additional information. As will be shown below, $\pi(D)$ is finite if and only if (\mathcal{G}_t) is a finite utility filtration.

We now approximate the general filtration (\mathcal{G}_t) by filtrations that can be seen as successive initial enlargements. Let $\Delta : 0 = s_0 \leq \dots \leq s_n = T$, $n \in \mathbb{N}$, be a partition of the interval $[0, T]$. We let for $r \in [s_i, s_{i+1})$, $i = 0, \dots, n-1$,

$$\mathcal{G}_r^\Delta = \mathcal{G}_{s_i-} \vee \mathcal{F}_r.$$

Proposition 4.3. *For $i = 0, \dots, n-1$, let μ^{s_i} be a \mathcal{G}^{s_i} -information drift and set $\mu_r^\Delta := \mu_r^{s_i}$ for $r \in [s_i, s_{i+1})$. Then μ^Δ is a \mathcal{G}_t^Δ -information drift. Moreover,*

$$\frac{1}{2} \int_0^T (\mu_r^\Delta)^2 d\langle M, M \rangle_r = \pi(D^\Delta),$$

where $D^\Delta := \{(s, t) \in D : \exists i \in \{0, \dots, n-1\} \text{ with } s < s_i \text{ and } t > s_i\}$.

Proof. It is straightforward to verify that μ^Δ is an information drift for (\mathcal{G}_t^Δ) . Moreover,

$$\begin{aligned} \frac{1}{2}E \int_0^T (\mu_r^\Delta)^2 d\langle M, M \rangle_r &= \frac{1}{2} \sum_{i=0}^{n-1} E \int_{s_i}^{s_{i+1}} (\mu_r^{s_i})^2 d\langle M, M \rangle_r \\ &= \frac{1}{2} \sum_{i=0}^{n-1} \left(E \int_{s_i}^T (\mu_r^{s_i})^2 d\langle M, M \rangle_r - E \int_{s_{i+1}}^T (\mu_r^{s_i})^2 d\langle M, M \rangle_r \right) \\ &= \sum_{i=0}^{n-1} \pi([0, s_i] \times (s_i, s_{i+1}]) = \pi(D^\Delta). \end{aligned}$$

□

We can now state the main theorem of this section.

Theorem 4.4. *Let Δ_n , $n \in \mathbb{N}$, be a sequence of partitions of the interval $[0, T]$ the mesh of which tends to 0. If $\pi(D)$ is finite, then the information drifts μ^{Δ_n} converge in $L^2(M)$ to a (\mathcal{G}_t) -information drift μ . Moreover, the utility gain of the insider satisfies*

$$u(\mathcal{G}_t, x) - u(\mathcal{F}_t, x) = \frac{1}{2}E \int_0^T \mu^2 d\langle M, M \rangle = \pi(D).$$

If $\pi(D)$ is infinite, then so is the utility gain of the insider.

The proof of the theorem is based on the following proposition.

Proposition 4.5. *If $\pi(D) < \infty$, then there exists a (\mathcal{G}_t) -information drift μ . Moreover,*

$$\frac{1}{2}\|\mu\|_{L^2(M)}^2 = \pi(D).$$

Proof. Let Δ_n , $n \in \mathbb{N}$, be as in the above theorem with the additional assumption that Δ_{n+1} is a refinement of Δ_n for all $n \in \mathbb{N}$. Then one has $\mathcal{G}_t^{\Delta_n} \subset \mathcal{G}_t^{\Delta_{n+1}}$ for any $t \in [0, T]$. By proposition 4.3, $\frac{1}{2}\|\mu^{\Delta_n}\|_{L^2(M)}^2 = \pi(D^{\Delta_n}) \leq \pi(D)$. Due to theorem 3.5 the information drifts μ^{Δ_n} converge to a $(\bigvee_{n \in \mathbb{N}} \mathcal{G}_t^{\Delta_n}) = (\mathcal{G}_t)$ -information drift μ in $L^2(M)$. Using monotone convergence we obtain that

$$\pi(D) = \lim_{n \rightarrow \infty} \pi(D^{\Delta_n}) = \lim_{n \rightarrow \infty} \frac{1}{2}\|\mu^{\Delta_n}\|_{L^2(M)}^2 = \frac{1}{2}\|\mu\|_{L^2(M)}^2.$$

Since every cadlag (\mathcal{G}_t) -martingale is as well a (\mathcal{G}_t) -martingale, μ is a (\mathcal{G}_t) -information drift. □

Proof of theorem 4.4. Assume that $\pi(D)$ is finite. Since the mesh of the partitions Δ_n tends to zero, one has $\lim_{n \rightarrow \infty} 1_{D^{\Delta_n}}(x) = 1$ for all $x \in D$. Consequently, the dominated convergence theorem yields

$$\lim_{n \rightarrow \infty} \pi(D^{\Delta_n}) = \pi(D). \quad (11)$$

We established the existence of a (\mathcal{G}_t) -information drift μ in Proposition 4.5. Recall that by Lemma 3.4, the processes μ^{Δ_n} and $\mu - \mu^{\Delta_n}$ are orthogonal in $L^2(M)$. Consequently,

$$\|\mu - \mu^{\Delta_n}\|_{L^2(M)}^2 = \|\mu\|_{L^2(M)}^2 - \|\mu^{\Delta_n}\|_{L^2(M)}^2.$$

Due to (11) the right hand side of the previous equation converges to 0. Hence, μ^{Δ_n} converges to μ in $L^2(M)$. The remaining statements are consequences of Proposition 4.5 and Proposition 1.4. □

5 Additional utility and entropy of filtrations

In this section we consider the link between the additional expected logarithmic utility of a better informed agent and the entropy of the additional information he possesses. For simple initially enlarged filtrations, this connection was discovered in Amendinger et al. [AIS98]. Here we shall see that the expected logarithmic utility increment is given by an integral version of relative entropies of the σ -algebras of the filtration. This notion can best be understood as the limit of discrete entropy sums along a sequence of partitions of the trading interval as the mesh goes to 0. Alternatively, we are able to give an interpretation of the utility increment by *Shannon information* differences between the filtrations of the agents. In particular, we shall see that this differences are independent of any local martingales the filtrations may carry.

Suppose that the assumptions of chapter 2 are satisfied. Moreover, we assume that M is a continuous local martingale satisfying the (PRP) relative to (\mathcal{F}_t) . In this case the regular conditional probabilities relative to the σ -fields \mathcal{F}_t^0 may be written

$$P_t(\cdot, A) = P(A) + \int_0^t k_s(\cdot, A) dM_s,$$

where $k(\cdot, A)$ is (\mathcal{F}_t) -predictable. Let again (\mathcal{G}_t^0) be a filtration satisfying $\mathcal{F}_t^0 \subset \mathcal{G}_t^0$ and being generated by countably many sets. To simplify notation we assume the filtration (\mathcal{G}_t^0) to be left-continuous. Let (\mathcal{G}_t) be the smallest completed and right-continuous filtration containing (\mathcal{G}_t^0) . In the following, we assume that (\mathcal{G}_t) is a finite utility filtration and denote by μ its predictable information drift, i.e.

$$\tilde{M} = M - \int_0^\cdot \mu_t d\langle M, M \rangle_t$$

is a (\mathcal{G}_t) -local martingale. Recall that by Theorem 2.10 we may assume that $k_t(\omega, \cdot)$ is a signed measure. For a fixed $r > 0$ we define μ^r as the information drift of the initially enlarged filtration (\mathcal{G}_t^r) , defined as in the beginning of the preceding chapter. For stating the main result we need the following Lemma.

Lemma 5.1. *Let $0 \leq s < t$ and $(\mathcal{P}^m)_{m \geq 0}$ an increasing sequence of finite partitions such that $\sigma(\mathcal{P}^m : m \geq 0) = \mathcal{G}_s^0$. Then*

$$\lim_m E \int_s^t \sum_{A \in \mathcal{P}^m} \left(\frac{k_u}{P_u} \right)^2 (\cdot, A) 1_A d\langle M, M \rangle_u = E \int_s^t (\mu_u^s)^2 d\langle M, M \rangle_u$$

and

$$\lim_m E \int_s^t \sum_{A \in \mathcal{P}^m} \frac{k_u}{P_u} (\cdot, A) 1_A \mu_u^s d\langle M, M \rangle_u = E \int_s^t (\mu_u^s)^2 d\langle M, M \rangle_u.$$

Proof. By Lemma 2.10 the process

$$Y_u^m(\omega, \omega') = \sum_{A \in \mathcal{P}^m} \frac{k_u}{P_u}(\omega, A) 1_A(\omega'), \quad m \geq 1,$$

is a L^2 -bounded martingale for P_M -a.a. $(\omega, u) \in \Omega \times [s, t]$. Hence (Y^m) converges P_M -a.s. to the density

$$\gamma_u = \frac{k_u(\cdot, d\omega')}{P_u(\cdot, d\omega')} \Big|_{\mathcal{G}_s^0}.$$

By Theorem 2.6 we have

$$\gamma_u(\omega, \omega') = \mu_u^s(\omega)$$

P_M -a.s. on $\Omega \times [s, t]$ and hence the first result. In a similar way one can prove the second statement. \square

We next discuss the important concept of the additional information of a σ -field relative to a filtration.

Definition 5.2. Let \mathcal{A} be a sub- σ -algebra of \mathcal{F} and R, Q two probability measures on \mathcal{F} . Then we define the relative entropy of R with respect to Q on the σ -field \mathcal{A} by

$$\mathcal{H}_{\mathcal{A}}(R\|Q) = \begin{cases} \int \log \frac{dR}{dQ} \Big|_{\mathcal{A}} dR, & \text{if } R \ll Q \\ \infty, & \text{else.} \end{cases}$$

Moreover, the *additional information* of \mathcal{A} relative to the filtration (\mathcal{F}_r) on $[s, t]$ ($0 \leq s < t \leq T$) is defined by

$$H_{\mathcal{A}}(s, t) = \int \mathcal{H}_{\mathcal{A}}(P_t(\omega, \cdot) \| P_s(\omega, \cdot)) dP(\omega).$$

The following lemma establishes the basic link between the entropy of a filtration enlargement and additional logarithmic utility of a trader possessing this information advantage.

Lemma 5.3. For $0 \leq s < t$ we have

$$H_{\mathcal{G}_s^0}(s, t) = \frac{1}{2} E \int_s^t (\mu_u^s)^2 d\langle M, M \rangle_u.$$

Proof. Let $(\mathcal{P}^m)_{m \geq 0}$ be an increasing sequence of finite partitions such that $\sigma(\mathcal{P}^m : m \geq 0) = \mathcal{G}_s^0$. Recall that by equation (8)

$$\begin{aligned} & \sum_{A \in \mathcal{P}^m} [1_A \log P_s(\cdot, A) - 1_A \log P_t(\cdot, A)] \\ = & \sum_{A \in \mathcal{P}^m} \left[- \int_s^t \frac{k_u}{P_u}(\cdot, A) 1_A d\tilde{M}_u - \int_s^t \frac{k_u}{P_u}(\cdot, A) 1_A \mu_u d\langle M, M \rangle_u \right. \\ & \left. + \frac{1}{2} \int_s^t \left(\frac{k_u}{P_u} \right)^2 (\cdot, A) 1_A d\langle M, M \rangle_u \right] \end{aligned}$$

Since \tilde{M} is a local martingale, we obtain by stopping and taking limits if necessary

$$\begin{aligned} & E \sum_{A \in \mathcal{P}^m} P_s(\cdot, A) \log \frac{P_t(\cdot, A)}{P_s(\cdot, A)} \\ = & E \sum_{A \in \mathcal{P}^m} \int_s^t \frac{k_u}{P_u}(\cdot, A) 1_A \mu_u d\langle M, M \rangle_u - \frac{1}{2} \int_s^t \left(\frac{k_u}{P_u} \right)^2 (\cdot, A) 1_A d\langle M, M \rangle_u. \end{aligned}$$

Note that in the previous line μ may be replaced by μ^s , because $(\mu - \mu^s)$ is orthogonal to $L^2(M)(\mathcal{G}^s)$ (see property iv) in Lemma 1.6). Applying Lemma 5.1 yields

$$\lim_m H_{\mathcal{P}^m}(s, t) = \frac{1}{2} E \int_s^t (\mu_u^s)^2 d\langle M, M \rangle_u.$$

Fatou's Lemma implies

$$\liminf_m H_{\mathcal{P}^m}(s, t) \geq H_{\mathcal{G}_s^0}(s, t).$$

On the other hand we have $H_{\mathcal{P}^m}(s, t) \leq H_{\mathcal{G}_s^0}(s, t)$, since $\mathcal{P}^m \subset \mathcal{G}_s^0$, and thus

$$\lim_m H_{\mathcal{P}^m}(s, t) = H_{\mathcal{G}_s^0}(s, t),$$

which completes the proof. \square

Let us now return to the stepwise approximation of a filtration enlargement along a sequence of partitions of the trading interval by "initial enlargements", and define their respective information increment.

Definition 5.4. Let $\Delta : 0 = s_0 \leq \dots \leq s_n = T$, $n \in \mathbb{N}$, be a partition of the interval $[0, T]$ and let μ^Δ be the information drift of (\mathcal{G}_r^Δ) . The additional information of (\mathcal{G}_r^Δ) relative to (\mathcal{F}_r) is defined as

$$H_\Delta = \sum_{i=0}^{n-1} H_{\mathcal{G}_{s_i}^0}(s_i, s_{i+1}).$$

Theorem 5.5. *We have*

$$\lim_{|\Delta| \rightarrow 0} H_\Delta = \frac{1}{2} E \int_0^T \mu_u^2 d\langle M, M \rangle_u.$$

Proof. This follows from Theorem 4.4 and Lemma 5.3. \square

Example 5.6. Let $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \sigma(\mathcal{P})$, where \mathcal{P} is a finite partition in \mathcal{F}_T . Then $\mu^0 = \mu$ and by Lemma 5.3

$$H_{\mathcal{G}_0^0}(0, T) = \frac{1}{2} E \int_0^T \mu_u^2 d\langle M, M \rangle_u.$$

If \mathcal{F}_0 is trivial, then

$$H_{\mathcal{G}_0^0}(0, T) = - \sum_{A \in \mathcal{P}} P(A) \log P(A),$$

which is the absolute entropy of the partition \mathcal{P} . Thus, the additional logarithmic utility of an agent with information (\mathcal{G}_t) is equal to the entropy of \mathcal{P} . This example shows that there is a link between logarithmic utility and the so-called Shannon information.

Definition 5.7. Let X and Y be two random variables in some measurable spaces. The *mutual information* (or Shannon information) between X and Y is defined by

$$I(X, Y) = \mathcal{H}(P_{X,Y} \| P_X \otimes P_Y).$$

Now let Z be a third random variable. The *conditional mutual information* of X and Y given Z is defined by

$$I(X, Y|Z) = \mathbb{E} \left[\mathcal{H}(P_{X,Y|Z} \| P_{X|Z} \otimes P_{Y|Z}) \right],$$

provided the regular conditional probabilities exist.

If \mathcal{A} is a sub- σ -algebra of \mathcal{F} , then we write $\text{id}_{\mathcal{A}}$ for the measurable map $(\Omega, \mathcal{F}) \rightarrow (\Omega, \mathcal{A})$, $\omega \mapsto \omega$. For two sub- σ -algebras \mathcal{A} and \mathcal{D} we abbreviate

$$I(\mathcal{A}, \mathcal{D}) = I(\text{id}_{\mathcal{A}}, \text{id}_{\mathcal{D}}).$$

Since our probability space is standard, for any sub- σ -fields $\mathcal{A}, \mathcal{D}, \mathcal{E}$ of \mathcal{F} there exists a regular conditional probability $P_{\text{id}_{\mathcal{A}}, \text{id}_{\mathcal{D}} | \text{id}_{\mathcal{E}}}$ and we define

$$I(\mathcal{A}, \mathcal{D} | \mathcal{E}) := I(\text{id}_{\mathcal{A}}, \text{id}_{\mathcal{D}} | \text{id}_{\mathcal{E}}).$$

The mutual information was introduced by Shannon as a measure of information. It plays an important role in information theory (see, for instance, [Iha93]).

Lemma 5.8. *For $0 \leq s < t \leq T$ we have*

$$H_{\mathcal{G}_s^0}(s, t) = I(\mathcal{G}_s^0, \mathcal{F}_t^0 | \mathcal{F}_s^0).$$

Proof. Note that for three random variables X, Y and Z we have

$$\frac{dP_{(X,Y)|Z}}{d(P_{X|Z} \otimes P_{Y|Z})} = \frac{dP_{X|(Y,Z)}}{dP_{X|Z}}.$$

This property implies

$$\begin{aligned} I(\mathcal{G}_s^0, \mathcal{F}_t^0 | \mathcal{F}_s^0) &= \int \int \log \frac{dP_{\text{id}_{\mathcal{G}_s^0} | \text{id}_{\mathcal{F}_t^0}}}{dP_{\text{id}_{\mathcal{G}_s^0} | \text{id}_{\mathcal{F}_s^0}}} dP(\omega') dP(\omega) \\ &= \int \int \log \frac{P_t(\cdot, d\omega')}{P_s(\cdot, d\omega')} \Big|_{\mathcal{G}_s^0} dP(\omega') dP(\omega) \\ &= H_{\mathcal{G}_s^0}(s, t). \end{aligned}$$

□

As a consequence we get

Theorem 5.9.

$$\lim_{|\Delta| \rightarrow 0} \sum_i I(\mathcal{G}_{s_i}^0, \mathcal{F}_{s_{i+1}}^0 | \mathcal{F}_{s_i}^0) = \frac{1}{2} E \int_0^T \mu_u^2 d\langle M, M \rangle_u.$$

Proof. Follows immediately from Theorem 5.5. □

This result motivates the following notion.

Definition 5.10. The *information difference* of (\mathcal{G}_r^0) relative to (\mathcal{F}_r^0) up to time T is defined as

$$\lim_{|\Delta| \rightarrow 0} \sum_i I(\mathcal{G}_{s_i}^0, \mathcal{F}_{s_{i+1}}^0 | \mathcal{F}_{s_i}^0).$$

We will denote this limit by $A(\mathcal{G}^0, \mathcal{F}^0)$.

Remark 5.11. Note that we did not use M in our definition of the information difference of (\mathcal{G}_r^0) relative to (\mathcal{F}_r^0) . However, by Theorem 5.9, the information difference may be represented in terms of any local martingale satisfying the (PRP).

Theorem 5.9 can be reformulated in the following way.

Theorem 5.12. *The additional utility of an agent with information (\mathcal{G}_t) is equal to the information difference of (\mathcal{G}_r^0) relative to (\mathcal{F}_r^0) , i.e.*

$$u_{\mathcal{G}}(x) - u_{\mathcal{F}}(x) = A(\mathcal{G}^0, \mathcal{F}^0).$$

If (\mathcal{G}_t) is initially enlarged by some random variable G , then the information difference of (\mathcal{G}_r^0) relative to (\mathcal{F}_r^0) coincides with the Shannon information between G and (\mathcal{F}_T^0) .

Lemma 5.13. *Let $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \sigma(G)$, where G is a random variable with values in some Polish space. Then*

$$A(\mathcal{G}^0, \mathcal{F}^0) = I(G, \mathcal{F}_T^0 | \mathcal{F}_0^0).$$

Proof. Let $0 \leq s \leq t$. By standard arguments we have

$$I(\mathcal{G}_s^0, \mathcal{F}_t^0 | \mathcal{F}_s^0) = I(G, \mathcal{F}_t^0 | \mathcal{F}_s^0)$$

and

$$\begin{aligned} I(G, \mathcal{F}_t^0 | \mathcal{F}_0^0) &= I(G, (\mathcal{F}_t^0, \mathcal{F}_s^0) | \mathcal{F}_0^0) \\ &= I(G, \mathcal{F}_t^0 | \mathcal{F}_s^0) + I(G, \mathcal{F}_s^0 | \mathcal{F}_0^0) \end{aligned}$$

(see e.g. [Iha93] Theorem 1.6.3.) By iteration we obtain for all partitions Δ

$$\sum_i I(\mathcal{G}_{s_i}^0, \mathcal{F}_{s_{i+1}}^0 | \mathcal{F}_{s_i}^0) = I(G, \mathcal{F}_T^0 | \mathcal{F}_0^0),$$

and hence the result. □

Theorem 5.14. *Let $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \sigma(G)$, where G is a random variable with values in some Polish space. Then the additional logarithmic utility of an agent with information (\mathcal{G}_t) is equal to the Shannon information between G and (\mathcal{F}_T^0) conditioned on \mathcal{F}_0 , i.e.*

$$u_{\mathcal{G}}(x) - u_{\mathcal{F}}(x) = I(\mathcal{F}_T^0, G | \mathcal{F}_0^0).$$

In particular, if \mathcal{F}_0^0 is trivial, then the additional utility is equal to $I(\mathcal{F}_T^0, G)$.

Proof. This follows from Lemma 5.13 and Theorem 5.9. □

Remark 5.15. If $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \sigma(G)$ and G is \mathcal{F}_T^0 -measurable, then the mutual information $I(\mathcal{F}_T^0, G | \mathcal{F}_0^0)$ is equal to the conditional absolute entropy of G (see also [AIS98]).

Example 5.16. Let (Ω, \mathcal{F}, P) be the 1-dimensional canonical Wiener space equipped with the Wiener process $(W_t)_{0 \leq t \leq 1}$. More precisely, $\Omega = \mathcal{C}([0, 1], \mathbb{R})$ is the set of continuous functions on $[0, 1]$ starting in 0, \mathcal{F} the σ -algebra of Borel sets with respect to uniform convergence, P the Wiener measure and W the coordinate process. $(\mathcal{F}_t)_{0 \leq t \leq 1}$ is obtained by completing the natural filtration $(\mathcal{F}_t^0)_{0 \leq t \leq 1}$. Suppose the price process S is of the form

$$S_t = \exp(W_t + bt), \quad 0 \leq t \leq 1,$$

with $b \in \mathbb{R}$. We want to calculate the additional utility of an insider knowing whether the price exceeds a certain level or not. More precisely, we suppose the insider to know the value of

$$G = 1_{(c, \infty)}(S_1^*),$$

where $c > 0$ and $S_1^* = \max_{0 \leq t \leq 1} S_t$. By Remark 5.15 the additional utility is equal to the entropy

$$H(G) = p \log p + (1 - p) \log(1 - p)$$

where

$$p = P(S_1^* > c).$$

This may be calculated via Girsanov's theorem. Namely we have

$$\begin{aligned} P(S_1^* > c) &= P(\forall t \in [0, 1] : W_t + bt > \log c) \\ &= \int_0^1 \exp\left(b \log c - \frac{b^2}{2}s\right) \frac{|\log c|}{\sqrt{2\pi s^3}} \exp\left(-\frac{x^2}{2s}\right) ds. \end{aligned}$$

6 Mutual information estimates

In this final section we apply some results from information theory to derive estimates for the information of a better informed agent. This yields a priori estimates for the agent's additional expected logarithmic utility in the light of the preceding section. Among other facts, the differential entropy maximizing property of Gaussian laws will play a role. We adopt the notations of [Iha93].

Before we provide the information estimates, we summarize some basic facts of the mutual information (see [Iha93], Theorem 1.6.3). For random variables X, Y, Z in some Borel spaces, the following properties hold:

(I.1) $I(X, Y|Z) \geq 0$ and, $I(X, Y|Z) = 0$ if and only if X and Y are independent given Z

(I.2) $I(X, (Y, Z)) = I(X, Z) + I(X, Y|Z)$

(I.3) If X is a continuous random variable with finite differential entropy, then

$$I(X, Y) = h(X) - h(X|Y).$$

For some fixed integer $d \in \mathbb{N}$, let X be a \mathcal{F}_T^0 -measurable \mathbb{R}^d -valued random variable. Moreover, denote by Y a d -dimensional r.v. that is independent of the σ -field \mathcal{F}_T^0 . We consider the enlarged filtration $\mathcal{G}_t^0 = \mathcal{F}_t^0 \vee \sigma(G)$, where $G := X + Y$.

Lemma 6.1. *Suppose that the law of Y is absolutely continuous with respect to Lebesgue measure and has finite differential entropy*

$$h(Y) = - \int \frac{dP_Y}{d\lambda^d}(y) \log \frac{dP_Y}{d\lambda^d}(y) dy.$$

Then

$$I(G, \mathcal{F}_T^0) = h(X + Y) - h(Y). \quad (12)$$

Proof. Due to property (I.2), we have

$$I(G, \mathcal{F}_T^0) = I(X + Y, X) + I(X + Y, \mathcal{F}_T^0|X).$$

Given X , the r.v.'s $X + Y$ and $\text{id}_{\mathcal{F}_T^0}$ are independent. Therefore, (I.1) and (I.3) lead to

$$I(G, \mathcal{F}_T^0) = I(X + Y, X) = h(X + Y) - h(X + Y|X) = h(X + Y) - h(Y).$$

□

Now assume the perturbation Y to be a \mathbb{R}^d -valued centered Gaussian r.v. that is independent of \mathcal{F}_T^0 .

Lemma 6.2. *Suppose that $X \in L^2(P)$ and let C_X and C_Y denote the covariance matrices of X and Y , respectively. Then*

$$I(G, \mathcal{F}_T^0) \leq \frac{1}{2} \log \frac{\det(C_X + C_Y)}{\det(C_Y)}. \quad (13)$$

Moreover, equality holds in equation (13) if X is Gaussian.

Proof. The distribution of Y is continuous with respect to Lebesgue measure and has finite entropy. Therefore,

$$I(G, \mathcal{F}_T^0) = h(X + Y) - h(Y).$$

Let C_X and C_Y denote the covariance matrices of X and Y , respectively. Due to the independence of X and Y , the random variable $X + Y$ has the covariance matrix $C_{X+Y} = C_X + C_Y$. Next recall that the normal distribution maximizes the differential entropy under a covariance constraint, i.e. $h(X + Y) \leq h(Z)$, where Z is a centered Gaussian r.v. with covariance matrix C_{X+Y} . Therefore,

$$I(X, X + Y) \leq h(Z) - h(Y).$$

Using the formula for the differential entropy of Gaussian measures (Theorem 1.8.1, [Iha93]) we obtain

$$\begin{aligned} h(Z) - h(Y) &= \frac{1}{2} \log((2\pi e)^d \det(C_{X+Y})) - \frac{1}{2} \log((2\pi e)^d \det(C_Y)) \\ &= \frac{1}{2} \log \frac{\det(C_{X+Y})}{\det(C_Y)}. \end{aligned}$$

If X is Gaussian, then $h(X + Y) = h(Z)$ and, hence, the second statement of the lemma follows. □

Corollary 6.3. *Assume that additionally to the assumptions of the above lemma, the equation $Y = \kappa N$ is valid, where N is a d -dimensional standard normal r.v. and $\kappa > 0$. Then*

$$I(G, \mathcal{F}_T^0) \leq \frac{1}{2} \sum_{j=1}^d \log \frac{\lambda_j + \kappa}{\kappa},$$

where λ_j ($j = 1, \dots, n$) denote the eigenvalues of C_X .

Proof. The proof follows easily by computing the determinants in Lemma 6.2. \square

The proof of Lemma 6.2 is based on the fact that Gaussian distributions maximize the differential entropy under a constraint on the covariance structure. Let us recall the construction of entropy maximizing measures under a linear constraint.

Lemma 6.4. *Let $E \subset \mathbb{R}^d$ be a measurable set, $c > 0$ and $g : E \rightarrow [0, \infty)$ a measurable map. Assume that there exist constants $Z, t \geq 0$, such that the measure ν defined by*

$$\frac{d\nu}{d\lambda^d}(x) = \frac{1}{Z} e^{-tg(x)},$$

is a probability measure satisfying $E^\nu[g] = c$. Then ν is the unique probability measure maximizing the differential entropy among all continuous probability measures μ on E satisfying $E^\mu[g] = c$.

The entropy maximization problem is equivalent to minimizing the relative entropy $\mathcal{H}(\cdot \| \lambda^d)$. Hence, the problem can be treated under more general constraints by using results of (Csiszár [Csi75], Theorem 3.1).

Proof. Let μ be a continuous probability measure on E with $E^\mu[g] = c$. Then

$$\begin{aligned} \mathcal{H}(\mu \| \nu) &= E^\mu \log \frac{d\mu}{d\nu} = E^\mu \log \frac{d\mu}{d\lambda^d} + E^\mu \log \frac{d\lambda^d}{d\nu} \\ &= -h(\mu) + \log Z + tE^\mu g = -h(\mu) + \log Z + tE^\nu g \\ &= -h(\mu) - E^\nu \log \frac{e^{tg}}{Z} = -h(\mu) + h(\nu). \end{aligned}$$

Since $\mathcal{H}(\mu \| \nu) \geq 0$ and $\mathcal{H}(\mu \| \nu) = 0$ iff $\mu = \nu$, ν is the unique maximizer of the differential entropy. \square

Remark 6.5. The above lemma can be used to derive similar results as obtained in Lemma 6.2. For instance, for $E := \mathbb{R}$ and $g(x) := |x|$ one obtains that the two-sided exponential distribution maximizes the differential entropy under the constraint $E^\mu g = c$ ($c > 0$). In particular, the measure ν with $\frac{d\nu}{d\lambda}(x) = (2c)^{-1} e^{-|x|/c}$ satisfies

$$E^\nu[g] = c \quad \text{and} \quad h(\nu) = 1 + \log(2c).$$

Now let X be a real-valued r.v. in $L^1(P)$. Moreover, let $\kappa_1 := E[|X - EX|]$ and Y be a two-sided exponential distribution with $E|Y| =: \kappa_2$. Then due to Lemma 6.1,

$$I(G, \mathcal{F}_T^0) \leq \log \left(\frac{\kappa_1 + \kappa_2}{\kappa_2} \right).$$

Example 6.6. We consider the classical stock market model with one asset. Let $(\mathcal{F}_t^0)_{t \in [0, T]}$ be a Brownian filtration generated by the Brownian motion $(B_t)_{t \in [0, T]}$ and denote by (\mathcal{F}_t) its completion. The stock price is modeled by the process

$$S_t = S_0 \exp\{B_t + bt\},$$

where $S_0 > 0$ is the deterministic stock price at time 0 and $b \in \mathbb{R}$. For some fixed times $t_1, \dots, t_d \in (0, T]$ ($d \in \mathbb{N}$), let $X := (B_{t_i})_{i=1, \dots, d}$. We suppose that the insider bases his investment on the filtration $\mathcal{G}_t = \bigcap_{s > t} \mathcal{F}_s \vee \sigma(G)$, where $G = X + \kappa N$ and N is a standard normal r.v. in \mathbb{R}^d that is independent of \mathcal{F}_T . Due to Lemma 6.2 the additional utility of the insider is related to the eigenvalues of the matrix

$$\begin{pmatrix} t_1 & t_1 & \dots & t_1 \\ t_1 & t_2 & \dots & t_2 \\ \vdots & \vdots & & \vdots \\ t_1 & t_2 & \dots & t_d \end{pmatrix}.$$

Let us finish the section with an example for a general enlargement.

Example 6.7. We reconsider the classical stock market model of Example 6.6 with $T := 1$. The knowledge of the insider at time t is modeled by $\mathcal{G}_t = \bigcap_{r > t} \mathcal{F}_r \vee \sigma((G_s)_{s \in [0, r]})$, where $G_t := B_1 + \tilde{B}_{g(1-t)}$, (\tilde{B}_t) is a Brownian motion independent of (B_t) and $g : [0, 1] \rightarrow [0, \infty)$ is a decreasing function. We are therefore in a setting similar to example 2.13. We now calculate the utility increment from the perspective of the notion of information difference of filtrations. Let π be as in section 4. For $0 \leq s \leq t \leq 1$ we have

$$\begin{aligned} \pi([0, s] \times (s, t]) &= I((G_u)_{u \in [0, s]}, \mathcal{F}_t^0 | \mathcal{F}_s^0) \\ &= I(G_s, \mathcal{F}_t^0 | \mathcal{F}_s^0) = I(G_s, B_t | \mathcal{F}_s^0) + I(G_s, \mathcal{F}_t^0 | \mathcal{F}_s^0, B_t) \\ &= I(B_1 + \tilde{B}_{g(1-s)}, B_t - B_s | \mathcal{F}_s^0) \\ &= I(B_1 - B_s + \tilde{B}_{g(1-s)}, B_t - B_s). \end{aligned}$$

Using property (I.8) we obtain

$$\begin{aligned} \pi([0, s] \times (s, t]) &= h(B_1 - B_s + \tilde{B}_{g(1-s)}) - h(B_1 - B_t + \tilde{B}_{g(1-s)}) \\ &= \frac{1}{2} \log(2\pi e(1-s+g(1-s))) - \frac{1}{2} \log(2\pi e(1-t+g(1-s))) \\ &= \frac{1}{2} \log \frac{1-s+g(1-s)}{1-t+g(1-s)} \end{aligned}$$

Alternatively one can express $\pi([0, s] \times (s, t])$ as

$$\pi([0, s] \times (s, t]) = \frac{1}{2} \int_s^t \frac{1}{1-u+g(1-s)} du.$$

For a partition $\Delta : 0 = t_0 \leq \dots \leq t_m = 1$ ($m \in \mathbb{N}$) we consider D^Δ as in section 4. One has

$$\begin{aligned} \pi(D^\Delta) &= \sum_{i=1}^n \pi([t_{i-1}, t_i] \times (t_i, t_{i+1}]) \\ &= \frac{1}{2} \int_0^1 \frac{1}{1-u+g(1-\max\{t_i : t_i \leq u\})} du \end{aligned}$$

Next, choose a sequence of refining partitions (Δ_n) such that their mesh tends to 0. Then the term in the latter integral is monotonically increasing in n and convergent. Hence, one obtains

$$\lim_{n \rightarrow \infty} \pi(D^{\Delta_n}) = \frac{1}{2} \int_0^1 \frac{1}{1-u+g(1-u)} du.$$

On the other hand,

$$\lim_{n \rightarrow \infty} \pi(D^{\Delta_n}) = \pi(D) = u_G(x) - u_{\mathcal{F}}(x).$$

Consequently the insider has finite utility if and only if $\int_0^1 \frac{1}{1-u+g(1-u)} du$. Now suppose $g(y) = Cy^p$ for some $C > 0$ and $p > 0$. It is straightforward to show that the integral, and hence the additional utility, is finite if and only if $p \in (0, 1)$. This equivalence follows also from results in [CIKHN03], where the authors compute explicitly the information drift.

References

- [ABS03] Jürgen Amendinger, Dirk Becherer, and Martin Schweizer. A monetary value for initial information in portfolio optimization. *Finance Stoch.*, 7(1):29–46, 2003.
- [AI04] Stefan Ankirchner and Peter Imkeller. Finite utility on financial markets with asymmetric information and the Theorem of Bichteler-Dellacherie-Mokobodzki. Preprint, 2004.
- [AIS98] Jürgen Amendinger, Peter Imkeller, and Martin Schweizer. Additional logarithmic utility of an insider. *Stochastic Process. Appl.*, 75(2):263–286, 1998.
- [Ank04] Stefan Ankirchner. Enlargement of filtrations and application to financial markets with asymmetric information. Ph.D. thesis, 2004.
- [Bau01] Fabrice Baudoin. Conditioning of brownian functionals and applications to the modelling of anticipations on a financial market. PhD thesis, Université Pierre et Marie Curie, 2001.
- [BO03] Francesca Biagini and Bernt Oksendal. A general stochastic calculus approach to insider trading. Preprint, 2003.
- [Cam03] Luigi Campi. Some results on quadratic hedging with insider trading. Preprint, Univ. Pescara, Univ. P. et M. Curie, 2003.
- [CIKHN03] Jose Corcuera, Peter Imkeller, Arturo Kohatsu-Higa, and David Nualart. Additional utility of insiders with imperfect dynamical information. *Preprint*, September 2003.
- [Csi75] Imre Csiszar. I-divergence geometry of probability distributions and minimization problems. *Ann. Probab.*, 3:146–158, 1975.
- [DH86] Darrell Duffie and Chi-fu Huang. Multiperiod security markets with differential information: martingales and resolution times. *J. Math. Econom.*, 15(3):283–303, 1986.

- [DM78] Claude Dellacherie and Paul-André Meyer. *Probabilities and potential*, volume 29 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1978.
- [DS95] Freddy Delbaen and Walter Schachermayer. The existence of absolutely continuous local martingale measures. *Ann. Appl. Probab.*, 5(4):926–945, 1995.
- [Els96] Jürgen Elstrodt. *Maß- und Integrationstheorie. (Measure and integration theory)*. Springer-Lehrbuch. Berlin: Springer, 1996.
- [GP98] Axel Grorud and Monique Pontier. Insider trading in a continuous time market model. *International Journal of Theoretical and Applied Finance*, 1:331–347, 1998.
- [GV03] Dario Gasbarra and Esko Valkeila. Initial enlargement: a Bayesian approach. Preprint, 2003.
- [Iha93] Shunsuke Ihara. *Information theory for continuous systems*. Singapore: World Scientific, 1993.
- [Imk96] Peter Imkeller. Enlargement of the Wiener filtration by an absolutely continuous random variable via Malliavin’s calculus. *Probab. Theory Related Fields*, 106(1):105–135, 1996.
- [Imk02] Peter Imkeller. Random times at which insiders can have free lunches. *Stochastics and Stochastics Reports*, 74:465–487, 2002.
- [Imk03] Peter Imkeller. Malliavin’s calculus in insider models: additional utility and free lunches. *Math. Finance*, 13(1):153–169, 2003. Conference on Applications of Malliavin Calculus in Finance (Rocquencourt, 2001).
- [IPW01] Peter Imkeller, Monique Pontier, and Ferenc Weisz. Free lunch and arbitrage possibilities in a financial market model with an insider. *Stochastic Process. Appl.*, 92(1):103–130, 2001.
- [JY85] Th. Jeulin and M. Yor, editors. *Grossissements de filtrations: exemples et applications*, volume 1118 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985. Papers from the seminar on stochastic calculus held at the Université de Paris VI, Paris, 1982/1983.
- [Par77] K.R. Parthasarathy. *Introduction to probability and measure*. Delhi etc.: MacMillan Co. of India Ltd. XII, , 1977.
- [PK96] Igor Pikovsky and Ioannis Karatzas. Anticipative portfolio optimization. *Adv. in Appl. Probab.*, 28(4):1095–1122, 1996.
- [RY99] Daniel Revuz and Marc Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999.