

TOPICS IN TOPOLOGY (“TOPOLOGIE III’), SOMMERSEMESTER 2024, HU BERLIN

CHRIS WENDL

This is not a set of lecture notes, but merely a brief summary of the contents of each lecture, with reading suggestions and a compendium of exercises. The suggested reading will usually not correspond precisely to what was covered in the lectures, but there will often be a heavy overlap.

PROLOGUE: NOTATION

Before getting into the content of the course, here is a glossary of important notation that is used in the lectures, including some comparison with other sources such as [tD08, DK01, Wen23] where different notation is sometimes used. This glossary will be updated during the semester as needed, and it is not in alphabetical order, but there is some kind of ordering principle... maybe you can figure out what it is.

Categories.

- General shorthand: For any category \mathcal{C} , I often abuse notation by writing $X \in \mathcal{C}$ to mean “ X is an **object** in \mathcal{C} ”; many other authors denote this by “ $X \in \text{Ob}(\mathcal{C})$ ” or something similar. For two objects $X, Y \in \mathcal{C}$, I write

$$\text{Hom}_{\mathcal{C}}(X, Y) \quad \text{or sometimes just} \quad \text{Hom}(X, Y)$$

for the set of **morphisms** $X \rightarrow Y$. The notation $\text{Mor}(X, Y)$ is also frequently used in many sources, and would make more sense linguistically, but it seems to be less popular. Given two functors $\mathcal{F}, \mathcal{G} : \mathcal{A} \rightarrow \mathcal{B}$, the notation

$$T : \mathcal{F} \rightarrow \mathcal{G}$$

means that T is a **natural transformation** from \mathcal{F} to \mathcal{G} .

- **Top**: the category of topological spaces and continuous maps
- **Top_{*}**: the category of **pointed spaces** and **pointed maps**, i.e. an object (X, x) is a topological space X equipped with a base point $x \in X$, and morphisms $f : (X, x) \rightarrow (Y, y)$ are continuous maps $X \rightarrow Y$ that send x to y . This notation is common but not universal, e.g. [tD08] uses a superscript 0 to indicate base points, so **Top_{*}** is called **TOP⁰**.
- **Set**: the category of sets and maps (with no continuity requirement)
- **Set_{*}**: the category of **pointed sets** and (not necessarily continuous) **pointed maps**, i.e. an object (X, x) is a set X with a base point $x \in X$, and morphisms $f : (X, x) \rightarrow (Y, y)$ are arbitrary maps $X \rightarrow Y$ that send x to y .
- **Top^{rel}**: the category of **pairs of spaces** (X, A) and **maps of pairs**, i.e. an object (X, A) is a topological space X equipped with a subset $A \subset X$, and morphisms $f : (X, A) \rightarrow (Y, B)$ are continuous maps $X \rightarrow Y$ that send A into B . Despite the unicity of this category, there doesn’t seem to be any common standard notation for it; [tD08] calls it **TOP(2)**, and similarly writes **TOP(3)** for the category of **triples** (X, A, B) with $B \subset A \subset X$, and so forth. In [Wen23] I used a subscript instead of a superscript, but I’m changing it so that I can also define the next item on this list.

- $\mathbf{Top}_*^{\text{rel}}$: the category of **pointed pairs of spaces**, i.e. an object (X, A, x) is a topological space X equipped with a subset $A \subset X$ and a base point $x \in A$, and morphisms $f : (X, A, x) \rightarrow (Y, B, y)$ are maps of pairs $(X, A) \rightarrow (Y, B)$ that also send x to y . I have no idea what anyone else calls this, but it's a subcategory of what [tD08] calls $\mathbf{TOP}(3)$, and is in any case clearly important since e.g. it is the domain of the relative homotopy functors π_n .
- \mathbf{hTop} , \mathbf{hTop}_* , $\mathbf{hTop}^{\text{rel}}$, $\mathbf{hTop}_*^{\text{rel}}$: the **homotopy categories** associated to \mathbf{Top} , \mathbf{Top}_* , $\mathbf{Top}^{\text{rel}}$ and $\mathbf{Top}_*^{\text{rel}}$ respectively, meaning we define categories with the same objects, but instead of taking morphisms to be actual maps, we define them to be *homotopy classes* of maps (respecting subsets and/or base points where appropriate, so e.g. **pointed homotopy** for \mathbf{hTop}_* , and homotopy of maps of pairs for $\mathbf{hTop}^{\text{rel}}$). This notation (or similar) for homotopy categories is very common, but different from my Topology I–II notes [Wen23], which wrote e.g. \mathbf{Top}_*^h instead of \mathbf{hTop}_* .
- \mathbf{Diff} : the category of smooth finite-dimensional **manifolds** without boundary, and **smooth maps**
- \mathbf{Grp} : the category of **groups** and group homomorphisms
- \mathbf{Ab} : the category of **abelian groups** and homomorphisms, which is a subcategory of \mathbf{Grp}
- $\mathbf{Ring} \supset \mathbf{CRing} \supset \mathbf{Fld}$: the category of **rings with unit** and its subcategories of **commutative rings** and **fields** respectively, with ring homomorphisms (preserving the unit)
- $\mathbf{R}\text{-Mod}$: the category of **modules** over a given commutative ring R and **R -module homomorphisms**. In [Wen23] I called this \mathbf{Mod}^R , and other variations such as $\mathbf{Mod}\text{-}R$ are also common.
- $\mathbb{K}\text{-Vect}$: the category of **vector spaces** over a given field \mathbb{K} and **\mathbb{K} -linear maps**, i.e. this is $\mathbf{R}\text{-Mod}$ in the special case where R is a field \mathbb{K} . In [Wen23] I called this $\mathbf{Vec}_{\mathbb{K}}$.
- Categories of **(co-)chain complexes**: given any *additive* category \mathcal{A} such as \mathbf{Ab} or $\mathbf{R}\text{-Mod}$,

$$\mathbf{Ch}(\mathcal{A}) \quad \text{or sometimes simply} \quad \mathbf{Ch}$$

denotes the category of chain complexes $\dots \rightarrow A_{n+1} \rightarrow A_n \rightarrow A_{n-1} \rightarrow \dots$ formed out of objects and morphisms in \mathcal{A} , with the morphisms of $\mathbf{Ch}(\mathcal{A})$ defined to be **chain maps**. There is a similar category $\mathbf{CoCh}(\mathcal{A})$ of cochain complexes $\dots \rightarrow A_{n-1} \rightarrow A_n \rightarrow A_{n+1} \rightarrow \dots$, though I am not really happy with this notation and I doubt that anyone else is either. In [Wen23] I denoted $\mathbf{Ch}(\mathbf{Ab})$, $\mathbf{CoCh}(\mathbf{Ab})$, $\mathbf{Ch}(\mathbf{R}\text{-Mod})$ and $\mathbf{CoCh}(\mathbf{R}\text{-Mod})$ by \mathbf{Chain} , $\mathbf{Cochain}$, \mathbf{Chain}^R and $\mathbf{Cochain}^R$ respectively. One sometimes sees a meaningless subscript such as $\mathbf{Ch}_\bullet(\mathcal{A})$ added, but there are also meaningful subscripts that define important subcategories such as e.g. $\mathbf{Ch}_{\geq 0}(\mathcal{A})$, the chain complexes that are trivial in all negative degrees.

- **Homotopy categories of chain complexes**: analogously to the homotopy categories of spaces, one can take the objects in $\mathbf{Ch}(\mathcal{A})$ and define morphisms to be chain homotopy classes of chain maps instead of actual chain maps. The internet seems quite insistent that I should call the resulting category

$$\mathbf{K}(\mathcal{A}) := \text{the (naive) homotopy category associated to } \mathbf{Ch}(\mathcal{A}),$$

even though I'd rather call it $\mathbf{hCh}(\mathcal{A})$, and in [Wen23] I wrote e.g. \mathbf{Chain}^h instead of $\mathbf{K}(\mathbf{Ab})$; on occasion I have even seen $\mathbf{Ho}(\mathcal{A})$ in place of $\mathbf{K}(\mathcal{A})$. I have no idea what notation to use for the homotopy category of cochain complexes. People who like derived categories will tell you that there are other things more deserving of the name “homotopy category of chain complexes,” and I added the word “naive” above in order to avoid getting into conversations about it with those people, which would be completely unnecessary for the purposes of the present course.

- \mathbf{Top}_B , $\mathbf{Top}_{B,*}$, \mathbf{hTop}_B and $\mathbf{hTop}_{B,*}$: Given a space B , these are the various categories of (unpointed or pointed) **spaces over B** with **maps over B** or homotopy classes thereof, as defined in Week 4, Lecture 6. The notation used in [tD08] for \mathbf{Top}_B and \mathbf{hTop}_B is not identical but sufficiently similar; I cannot find definitions in [tD08] for the pointed variants $\mathbf{Top}_{B,*}$ and $\mathbf{hTop}_{B,*}$.

Topological constructions.

- $X \amalg Y$: This is how I write the **disjoint union** of two topological spaces (and similarly for pairs of spaces), and most sensible people use either this notation or $X \sqcup Y$, but [tD08] instead writes $X + Y$ and calls it the **topological sum** of X and Y , presumably because—like the direct sum of abelian groups and many other constructions that use the word “sum”—it is a coproduct. The book by tom Tieck becomes significantly easier to read once you realize this.
- $X \coprod Y$: the **coproduct** of X and Y , whatever that means in whichever category X and Y happen to live in, so e.g. in \mathbf{Top} , it means the same thing as $X \amalg Y$, though in \mathbf{Top}_* it means $X \vee Y$.
- $[X, Y]$: If X and Y are just topological spaces (i.e. objects in \mathbf{Top}), then this denotes the **set of homotopy classes** of maps $X \rightarrow Y$, i.e.

$$[X, Y] := \mathrm{Hom}_{\mathbf{hTop}}(X, Y).$$

If X and Y are equipped with additional data (which may be suppressed in the notation) and are thus objects in \mathbf{Top}_* , $\mathbf{Top}^{\mathrm{rel}}$ or $\mathbf{Top}_*^{\mathrm{rel}}$, then I use the same notation $[X, Y]$ to mean the corresponding notion of homotopy classes in each category, so e.g. in the context of pointed spaces, I would write

$$[X, Y] := \mathrm{Hom}_{\mathbf{hTop}_*}(X, Y),$$

and similarly for (pointed or unpointed) pairs of spaces. This convention is popular but not universal, e.g. [tD08] writes $[X, Y]^0$ for the set of pointed homotopy classes and uses $[X, Y]$ only to mean unpointed homotopy classes; [DK01] does the same but writes $[X, Y]_0$ instead of $[X, Y]^0$.

- $X \vee Y$ and $X \wedge Y$: these are the **wedge sum** and **smash product** respectively of pointed spaces, and mercifully, everyone seems to agree on what they mean and how to write them.
- **Implied base points**: for a pair of spaces (X, A) , the **quotient space** X/A is often interpreted as a pointed space, with the collapsed subset A as base point. Similarly, for two pointed spaces X, Y , the set of **pointed homotopy classes** $[X, Y]$ is viewed as a pointed set (i.e. an object in \mathbf{Set}_*) whose base point is the homotopy class of the constant map to the base point of Y .
- **One-point spaces**: the symbol $*$ is often used to mean either a one-point space, the unique point in that space, or sometimes a previously unnamed base point of a given pointed space. It should usually be clear from context which is meant.
- I : this usually denotes the **unit interval**

$$I := [0, 1],$$

as appears in domains of paths, homotopies etc.

- **Homotopy relations**: Given maps $f, g : X \rightarrow Y$, I write

$$f \underset{h}{\sim} g$$

to mean that f and g are homotopic ([tD08] writes “ $f \simeq g$ ”), and

$$f \overset{H}{\rightsquigarrow} g$$

to mean that H is a homotopy from f to g , thought of as a *path* in the space of maps, hence $H : I \times X \rightarrow Y$ with $H(0, \cdot) = f$ and $H(1, \cdot) = g$. This can also mean e.g. pointed homotopy or homotopy of maps of pairs if working in \mathbf{Top}_* or $\mathbf{Top}^{\text{rel}}$ respectively. Where I write $f \xrightarrow{H} g$, [tD08] writes $H : f \simeq g$.

- **Homotopy commutative diagrams:** I use a diagram of the form

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ \downarrow g & \sim & \downarrow \varphi \\ Y & \xrightarrow{\psi} & Q \end{array}$$

to mean that $\varphi \circ f$ and $\psi \circ g$ need not be identical but are homotopic, whatever that means in whichever category the objects of the diagram live in, e.g. if they are pointed spaces it means pointed homotopic, for spaces without base points it just means homotopic—it may also mean *chain* homotopic if the objects are chain complexes. If I write the variant

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ \downarrow g & \underset{\sim}{\alpha} & \downarrow \varphi, \\ Y & \xrightarrow{\psi} & Q \end{array}$$

then it means that α is a homotopy (or chain homotopy as the case may be) from $\varphi \circ f$ to $\psi \circ g$. It wasn't easy to figure out how to render this in LaTeX, so maybe that's why most textbooks don't do it.

- $Z(f)$, $Z(f, g)$, $\text{cone}(f)$: **mapping cylinders**, **double mapping cylinders** and **mapping cones** (see Week 2, Lecture 3)
- CX , SX : the **cone** and **suspension** respectively of a space X . In the context of pointed spaces the same notation may instead mean the *reduced* cone/suspension.
- $P(f)$, $P(f, g)$, $F(f)$: the **mapping path space** of a map, **double mapping path space** of two maps, and **homotopy fiber** of a map respectively, as defined in Week 3, Lecture 5. These constructions are dual to $Z(f)$, $Z(f, g)$ and $\text{cone}(f)$ respectively, in the sense that they fit into analogous diagrams with all arrows reversed.
- $[X, Y]_B$: For two spaces X, Y over another space B , this is the set of (unpointed or pointed) **homotopy classes of maps over B** , i.e. morphisms in the category \mathbf{hTop}_B or $\mathbf{hTop}_{B,*}$.

1. WEEK 1

Lecture 1 (15.04.2024): Motivation and colimits.

- Motivational theorem on exotic spheres (Milnor 1956): There exists a smooth manifold Y that is homeomorphic but not diffeomorphic to S^7 . (In fact, Kervaire and Milnor proved shortly afterwards that there are exactly 28 such manifolds up to diffeomorphism.)
- Outline of a proof (slightly ahistorical), with notions that will be major topics in this course written in red:
 - (1) **Pontryagin classes:** Associate topological invariants $p_k(E) \in H^{4k}(X; \mathbb{Z})$ for each $k \in \mathbb{N}$ to every isomorphism class of **vector bundles** E over a given space X . Since every smooth manifold M has a **tangent bundle** TM , we can define $p_k(M) := p_k(TM) \in H^{4k}(M; \mathbb{Z})$ as an invariant of smooth (but not topological) manifolds.
 - (2) **Intersection form** and **signature:** For a compact oriented $4k$ -manifold M (possibly with boundary), the intersection form is the quadratic form Q_M on $H^{2k}(M, \partial M; \mathbb{Z})$ defined by

$$Q_M(\alpha) := \langle \alpha \cup \alpha, [M] \rangle \in \mathbb{Z},$$

and it’s called the “intersection form” because it can be interpreted as a signed count of intersections between two generic closed oriented submanifolds representing the class in $H_{2k}(M; \mathbb{Z})$ **Poincaré dual** to α . The signature $\sigma(M) \in \mathbb{Z}$ is essentially the number of positive eigenvalues minus the number of negative eigenvalues¹ of this quadratic form.

- (3) **Hirzebruch signature theorem** (8-dimensional case): For M a closed oriented 8-manifold,

$$\sigma(M) = \frac{1}{45} \langle 7p_2(M) - p_1(M) \cup p_1(M), [M] \rangle.$$

- (4) (the clever bit) Construct a compact oriented smooth 8-manifold X with simply connected boundary $Y := \partial X$ such that $\sigma(X) = 8$, $H_2(Y)$ and $H_3(Y)$ both vanish, and the tangent bundle TX is **stably trivial**, which implies its Pontryagin classes vanish. The construction can be described (key words: “plumbing of spheres”), and the computations carried out, using only methods from Topology 2.
- (5) Deduce via **Poincaré duality**, the **Hurewicz theorem** and **Whitehead’s theorem**² that Y is homotopy equivalent to S^7 . By Smale’s solution to the higher-dimensional Poincaré conjecture,³ it follows that Y is homeomorphic to S^7 .
- (6) Argue by contradiction: If Y is diffeomorphic to S^7 , then one can construct a closed smooth 8-manifold M by gluing X to an 8-disk along a diffeomorphism $\partial X = Y \cong S^7 = \partial \mathbb{D}^8$,

$$M := X \cup_{S^7} \mathbb{D}^8.$$

Methods from Topology 2 (e.g. Mayer-Vietoris) now imply $p_1(M) = 0$ and $\sigma(M) = 8$, so Hirzebruch says

$$45\sigma(M) = 45 \cdot 8 = 7 \langle p_2(M), [M] \rangle.$$

But the right hand side of this relation is a multiple of 7, and the left hand side is not.

- Interpretation of a functor $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$ as a **diagram** in \mathcal{C} over \mathcal{J} , constant functors $\mathcal{X} : \mathcal{J} \rightarrow \mathcal{C}$ as **targets**, the **universal property** and definition of the **colimit** $\text{colim}(\mathcal{F})$
- Interpreting direct systems as diagrams and direct limits as colimits
- Defining the quotient space X/A as colimit of the diagram

$$\begin{array}{ccc} A & \longrightarrow & * \\ \downarrow & & \\ X & & \end{array}$$

understood as a functor $\mathcal{J} \rightarrow \mathbf{Top}$, where \mathcal{J} is a category with three objects and only two nontrivial morphisms.

¹What I really mean is: first rewrite Q_M as a quadratic form on $H^{2k}(M, \partial M; \mathbb{Q})$ or $H^{2k}(M, \partial M; \mathbb{R})$, which is a vector space, so that by standard linear algebra, you can present it in terms of a symmetric linear transformation and look at the eigenvalues of that transformation. One can define this in a more obviously invariant way by talking about maximal subspaces on which Q_M is positive/negative definite.

²A 3-dimensional version of this same argument is described in [Wen23, Lecture 57], using the theorems of Hurewicz and Whitehead as black boxes.

³This is the one major black box in this proof that I do not intend to fill in, because that would be a whole course in itself.

Lecture 2 (18.04.2024): From coproducts to pullbacks and pushouts.

- The **limit** $\lim(\mathcal{F})$ of a diagram $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$
- Inverse limits as limits of diagrams
- Important special cases of limits and colimits:
 - **Coproducts** \coprod , and examples in the categories **Top** (disjoint union), **Top_{*}** (wedge sum), **Ab** (direct sum) and **Grp** (free product)
 - **Products** \times (or \prod), and examples in **Top**
 - **Equalizers** and **co-equalizers**, realization in **Top** as subspaces or quotient spaces respectively
- Word of caution: limits and colimits are not guaranteed to exist, e.g. in the category **Diff** of smooth finite-dimensional manifolds without boundary, finite or countable coproducts exist (and are the same thing as in **Top**), but uncountable disjoint unions are not second countable and are thus not objects in **Diff**. Similarly, finite products exist in **Diff** but infinite products typically do not.
- Theorem: In any category \mathcal{C} , all (co-)limits can be presented in terms of (co-)products and (co-)equalizers, if they exist.
- Proof sketch (co-limit case): Given $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C} : \alpha \mapsto X_\alpha$, construct $\text{colim}(\mathcal{F})$ as the equalizer of two morphisms $Y \xrightarrow{f,g} Z$ defined as follows. Write the set of all morphisms in \mathcal{J} as $\text{Hom}(\mathcal{J}, \mathcal{J})$; we then take Y to be the coproduct

$$Y := \coprod_{\phi \in \text{Hom}(\mathcal{J}, \mathcal{J})} X_\phi, \quad \text{where} \quad \text{for } \phi \in \text{Hom}(\alpha, \beta), X_\phi := X_\alpha,$$

while Z is the slightly simpler coproduct

$$Z := \coprod_{\beta \in \mathcal{J}} X_\beta.$$

For each $\alpha, \beta \in \mathcal{J}$ and $\phi \in \text{Hom}(\alpha, \beta)$, let $f_\phi : X_\phi \rightarrow Z$ denote the composition of the morphism $\phi_* : X_\phi = X_\alpha \rightarrow X_\beta$ with the canonical morphism $X_\beta \rightarrow \coprod_{\gamma \in \mathcal{J}} X_\gamma$ of the coproduct; the universal property of the coproduct then dictates that the collection of morphisms $f_\phi : X_\phi \rightarrow Z$ determines a morphism $f : Y \rightarrow Z$. Similarly, $g : Y \rightarrow Z$ is determined by the collection of morphisms $g_\phi : X_\phi \rightarrow Z$ defined for each $\phi \in \text{Hom}(\alpha, \beta)$ as the compositions of $\text{Id}_{X_\alpha} : X_\phi = X_\alpha \rightarrow X_\alpha$ with the canonical morphism $X_\alpha \rightarrow \coprod_{\gamma \in \mathcal{J}} X_\gamma$. Now check that the universal property is satisfied (exercise).

- Upshot: In **Top**, colimits are quotients of disjoint unions, limits are subspaces of products.
- **Fiber products**: presenting the fiber product of two maps $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ in **Top** as the “intersection locus”

$$X \times_{f \times g} Y := \{(x, y) \in X \times Y \mid f(x) = g(y)\}$$

with the obvious projections to X and Y .

- Interpreting fiber products as **pullbacks**
- **Pushouts**: presenting the pushout of two maps $f : Z \rightarrow X$ and $g : Z \rightarrow Y$ in **Top** as “gluing spaces together” along a map:

$$X \cup_f Y := (X \amalg Y) / f(z) \sim g(z) \text{ for all } z \in Z.$$

- Question for thought: How many of these constructions of limits or colimits work in the homotopy categories **hTop** or **hTop_{*}**? (Hint: Do not try too hard to make sense of equalizers and co-equalizers.)

Suggested reading. The main definitions involving direct systems and direct limits can all be found in [Wen23, Lecture 39], with the generalization to colimits explained in Exercise 39.24. If you're really serious about this stuff, you can also try reading [Mac71].

If you want to read more about exotic spheres, there's a nice collection of relevant literature assembled at <https://www.maths.ed.ac.uk/~v1ranick/exotic.htm>.

Exercises (for the Übung on 25.04.2024). Since the Übung on 25.04 was cancelled due to illness, most of the exercises for Week 1 have now been supplemented with written answers and/or some discussion.

Exercise 1.1. In what sense precisely are the limit and colimit of a diagram $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$ unique, if they exist?

Answer: If the limit or colimit exists (of which there is no guarantee, cf. Exercise 1.7), then it is unique up to canonical isomorphisms. Precisely: Suppose $X, Y \in \mathcal{C}$ are two objects, together with collections of morphisms $\mathcal{F}(\alpha) \xrightarrow{\varphi_\alpha} X$ and $\mathcal{F}(\alpha) \xrightarrow{\psi_\alpha} Y$ for all $\alpha \in \mathcal{J}$, such that both satisfy the universal property for $\text{colim}(\mathcal{F})$. Then there is a uniquely determined isomorphism

$$f : X \xrightarrow{\cong} Y \quad \text{such that} \quad \psi_\alpha = f \circ \varphi_\alpha \text{ for all } \alpha \in \mathcal{J}.$$

The existence and uniqueness of a morphism f satisfying this condition follows from the universal property of X , and the fact that it is an isomorphism follows by reversing the roles of X and Y , since Y also satisfies the universal property. For $\text{lim}(\mathcal{F})$ there is a similar uniqueness statement, proved in a similar way.

Note that in most categories, uniqueness "up to canonical isomorphisms" is the best that one could hope to get from universal properties, as one will always have the freedom to replace a given object playing the role of $\text{colim}(\mathcal{F})$ or $\text{lim}(\mathcal{F})$ with a different object that is isomorphic to it. In practice, our favorite categories often come with canonical constructions that lead to specific objects, e.g. the disjoint union (also known as the coproduct) of a given collection of topological spaces is a specific space, not just an equivalence class of spaces up to homeomorphism. But in various situations, limits or colimits can also arise from something other than the canonical construction, and finding an isomorphism with that canonical construction may be harder than explicitly verifying the universal property.

Exercise 1.2 (morphisms between (co-)products). Assume J is a set, and $\{X_\alpha\}_{\alpha \in J}$ and $\{Y_\alpha\}_{\alpha \in J}$ are collections of objects in some category \mathcal{C} such that the products

$$\left\{ \prod_{\alpha \in J} X_\alpha \xrightarrow{\pi_\beta^X} X_\beta \right\}_{\beta \in J}, \quad \left\{ \prod_{\alpha \in J} Y_\alpha \xrightarrow{\pi_\beta^Y} Y_\beta \right\}_{\beta \in J},$$

and coproducts

$$\left\{ X_\beta \xrightarrow{i_\beta^X} \coprod_{\alpha \in J} X_\alpha \right\}_{\beta \in J}, \quad \left\{ Y_\beta \xrightarrow{i_\beta^Y} \coprod_{\alpha \in J} Y_\alpha \right\}_{\beta \in J}$$

exist. In what sense does an arbitrary collection of morphisms $\{f_\alpha : X_\alpha \rightarrow Y_\alpha\}_{\alpha \in J}$ uniquely determine morphisms

$$\prod_{\alpha \in J} f_\alpha : \prod_{\alpha \in J} X_\alpha \rightarrow \prod_{\alpha \in J} Y_\alpha, \quad \text{and} \quad \coprod_{\alpha \in J} f_\alpha : \coprod_{\alpha \in J} X_\alpha \rightarrow \coprod_{\alpha \in J} Y_\alpha?$$

Argue in terms of universal properties, without using your knowledge of how to represent products and coproducts in any specific categories.

Answer: The morphisms $\coprod_{\alpha} f_{\alpha}$ and $\prod_{\alpha} f_{\alpha}$ are uniquely determined by the condition that the diagrams

$$\begin{array}{ccc} X_{\beta} & \xrightarrow{f_{\beta}} & Y_{\beta} \\ \downarrow i_{\beta}^X & & \downarrow i_{\beta}^Y \\ \coprod_{\alpha} X_{\alpha} & \xrightarrow{\coprod_{\alpha} f_{\alpha}} & \coprod_{\alpha} Y_{\alpha} \end{array} \quad \text{and} \quad \begin{array}{ccc} \prod_{\alpha} X_{\alpha} & \xrightarrow{\prod_{\alpha} f_{\alpha}} & \prod_{\alpha} Y_{\alpha} \\ \downarrow \pi_{\beta}^X & & \downarrow \pi_{\beta}^Y \\ X_{\beta} & \xrightarrow{f_{\beta}} & Y_{\beta} \end{array}$$

commute for every $\beta \in J$. One gets the existence and uniqueness of $\coprod_{\alpha} f_{\alpha}$ from the universal property of the coproduct $\coprod_{\alpha} X_{\alpha}$, because the morphisms $i_{\beta}^Y \circ f_{\beta} : X_{\beta} \rightarrow \coprod_{\alpha} Y_{\alpha}$ make $\coprod_{\alpha} Y_{\alpha}$ a target of the diagram whose colimit is $\coprod_{\alpha} X_{\alpha}$. Similarly, the existence and uniqueness of $\prod_{\alpha} f_{\alpha}$ follows from the universal property of the product $\prod_{\alpha} Y_{\alpha}$, using the collection morphisms $f_{\beta} \circ \pi_{\beta}^X : \prod_{\alpha} X_{\alpha} \rightarrow Y_{\beta}$.

Exercise 1.3 (finite limits and colimits). Show that in any category \mathcal{C} , finite colimits always exist if and only if all pushouts exist and \mathcal{C} has an initial object (see Exercise 1.5). Dually, finite limits always exist if and only if all pullbacks (also known as fiber products) exist and \mathcal{C} has a terminal object.⁴

Hint: By a theorem stated in the lecture, it suffices if you can express arbitrary (co-)equalizers and finite (co-)products in terms of pushouts or pullbacks.

Solution: Note that the statement of this exercise has been revised; the original version had two errors, one being its failure to mention initial and terminal objects, and the other an oversimplification of what it means for a limit or colimit to be *finite*—we need the category \mathcal{J} underlying the diagram to have finitely-many morphisms, not just finitely-many objects.

With that understood, let's assume all pushouts exist and that \mathcal{C} also has an initial object $0 \in \mathcal{C}$. If we can show that all finite coproducts and all coequalizers exist, then the theorem from lecture uses these to construct a colimit for any diagram $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$ such that \mathcal{J} has only finitely many objects and morphisms. (Regarding the errors in the original version: note that if \mathcal{J} has finitely-many objects but infinitely-many morphisms, then one of the coproducts needed for the theorem from lecture is not finite.)

You should be able to convince yourself via an inductive argument that if the coproduct of two objects $X, Y \in \mathcal{C}$ always exists, then all finite coproducts exist. So let's show first that $X \coprod Y$ exists for arbitrary $X, Y \in \mathcal{C}$. At this point I find it helpful to think about how coproducts and pushouts are constructed concretely in the example $\mathcal{C} = \mathbf{Top}$: the coproduct of X and Y is their disjoint union, and the pushout of a pair of maps $f : Z \rightarrow X$ and $g : Z \rightarrow Y$ is a quotient of that disjoint union by the equivalence relation such that $f(z) \sim g(z)$ for all $z \in Z$. If we want to make that equivalence relation trivial so that the pushout turns out to be the same thing as the coproduct, the solution is to choose the empty set for Z ; the maps f, g are uniquely determined by this choice, because the empty set is an initial object in \mathbf{Top} (see Exercise 1.5). This suggests that in our given category \mathcal{C} with initial object $0 \in \mathcal{C}$, the pushout of the diagram

$$\begin{array}{ccc} 0 & \longrightarrow & X \\ \downarrow & & \\ & & Y \end{array}$$

should be the coproduct of X and Y ; note that only one diagram of this form is possible since 0 being initial means that the morphisms $0 \rightarrow X$ and $0 \rightarrow Y$ are unique. Now suppose P is the

⁴The word “finite” in this context refers to limits or colimits of diagrams $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$ such that \mathcal{J} has only finitely many objects and morphisms.

pushout of this diagram, equipped with morphisms $\varphi : X \rightarrow P$ and $\psi : Y \rightarrow P$, and suppose we are given another object Z with morphisms $f : X \rightarrow Z$ and $g : Y \rightarrow Z$. The diagram

$$\begin{array}{ccc} 0 & \longrightarrow & X \\ \downarrow & & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

then trivially commutes, since there is only one morphism $0 \rightarrow Z$, and the universal property of the pushout gives rise to a unique morphism $u : P \rightarrow Z$ such that $f = u \circ \varphi$ and $g = u \circ \psi$, which amounts to the statement that P with its morphisms φ and ψ also satisfies the universal property of the coproduct $X \coprod Y$.

We show next that the coequalizer of an arbitrary pair of morphisms

$$X \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} Y$$

in \mathcal{C} can also be constructed as a pushout. Think again about how it works in the case $\mathcal{C} = \mathbf{Top}$: the coequalizer here is the quotient of Y by the equivalence relation such that $f(x) \sim g(x)$ for all $x \in X$. If we instead take the pushout of f and g , the resulting space is too large: it is a quotient of $Y \amalg Y$ instead of Y , meaning that we glue together *two* copies of Y by identifying $f(x)$ in one copy with $g(x)$ in the *other* copy for each $x \in X$. But the correct space can be obtained from this by making the equivalence relation larger, so that for every $y \in Y$, y in the first copy gets identified with y in the second copy. The way to realize this is by enlarging the domain of the pair of maps used in defining the pushout: instead of the two maps $f, g : X \rightarrow Y$, we consider the pushout of the two maps $f \amalg \text{Id}, g \amalg \text{Id} : X \amalg Y \rightarrow Y$.

Let's say that again without assuming $\mathcal{C} = \mathbf{Top}$. We've already shown that the coproduct $X \coprod Y$ of two objects in \mathcal{C} can be constructed, and if we write $i_X : X \rightarrow X \coprod Y$ and $i_Y : Y \rightarrow X \coprod Y$ for the canonical morphisms that coproducts come equipped with, then by the universal property of the coproduct, every morphism $\varphi : X \rightarrow Y$ determines a unique morphism $\varphi \amalg \text{Id} : X \coprod Y \rightarrow Y$ for which the diagram

$$\begin{array}{ccc} X & & \xrightarrow{\varphi} Y \\ \downarrow i_X & & \searrow \varphi \amalg \text{Id} \\ X \coprod Y & \xrightarrow{\varphi \amalg \text{Id}} & Y \\ \uparrow i_Y & & \nearrow \text{Id} \\ Y & & \end{array}$$

commutes. Claim: Given two morphisms $f, g : X \rightarrow Y$, a diagram of the form

$$\begin{array}{ccc} X \coprod Y & \xrightarrow{f \amalg \text{Id}} & Y \\ \downarrow g \amalg \text{Id} & & \downarrow \varphi \\ Y & \xrightarrow{\psi} & Z \end{array}$$

commutes if and only if $\varphi = \psi$ and $\varphi \circ f = \varphi \circ g$. To see this, we can enhance the diagram in two ways using the universal property of the coproduct: first,

$$\begin{array}{ccccc}
 & & \text{Id} & & \\
 & \curvearrowright & & \curvearrowleft & \\
 Y & & & & Y \\
 \downarrow i_Y & & & & \downarrow \varphi \\
 & X \amalg Y & \xrightarrow{f \amalg \text{Id}} & & Y \\
 \downarrow g \amalg \text{Id} & & & & \downarrow \varphi \\
 & Y & \xrightarrow{\psi} & & Z
 \end{array}$$

(Note: The diagram above is a simplified representation of the one in the image. The original image shows a more complex diagram with additional arrows from Y to X and Y to Y.)

shows that if the given diagram commutes, then $\varphi = \varphi \circ \text{Id} = \psi \circ \text{Id} = \psi$. Assuming this, the second enhanced diagram

$$\begin{array}{ccccc}
 & & f & & \\
 & \curvearrowright & & \curvearrowleft & \\
 X & & & & Y \\
 \downarrow i_X & & & & \downarrow \varphi \\
 & X \amalg Y & \xrightarrow{f \amalg \text{Id}} & & Y \\
 \downarrow g \amalg \text{Id} & & & & \downarrow \varphi \\
 & Y & \xrightarrow{\psi} & & Z
 \end{array}$$

(Note: The diagram above is a simplified representation of the one in the image. The original image shows a more complex diagram with additional arrows from X to X and X to Y.)

then proves $\varphi \circ f = \psi \circ g = \varphi \circ g$. Conversely, if one assumes $\varphi = \psi$ and $\varphi \circ f = \varphi \circ g$, then $\varphi \circ (f \amalg \text{Id})$ and $\psi \circ (g \amalg \text{Id})$ are two morphisms $X \amalg Y \rightarrow Z$ whose compositions with i_X and i_Y are identical, so the uniqueness in the universal property of the coproduct requires them to be the same.

The result of the claim is that pushout diagrams for the two morphisms $f \amalg \text{Id} : X \amalg Y \rightarrow Y$ and $g \amalg \text{Id} : X \amalg Y \rightarrow Y$ are equivalent to coequalizer diagrams for $f, g : X \rightarrow Y$. It is a short step from there to the conclusion that an object Z with morphism $Y \rightarrow Z$ satisfies the universal property of the coequalizer if and only if Z with two copies of that same morphism $Y \rightarrow Z$ satisfies the universal property of the pushout.

For the dual case of this whole story, I will just say this: if $1 \in \mathcal{C}$ is a terminal object, then the uniqueness of morphisms to 1 implies that the pullback of the diagram

$$\begin{array}{ccc}
 & X & \\
 & \downarrow & \\
 Y & \longrightarrow & 1
 \end{array}$$

satisfies the universal property of the product $X \times Y$. Having shown that finite products exist, one then obtains the equalizer of any pair of morphisms $f, g : X \rightarrow Y$ as the pullback of the diagram

$$\begin{array}{ccc}
 & X & \\
 & \downarrow \text{Id} \times f & \\
 X & \xrightarrow{\text{Id} \times g} & X \times Y
 \end{array}$$

If finite products and equalizers always exist, then all finite limits can be constructed out of them.

Exercise 1.4. Let's talk about some coproducts and products in algebraic settings.

- What is a coproduct of two objects in the category **Ring** of rings with unit? Try to describe it explicitly.
- Same question about products in **Ring**. (This one is perhaps easier.)
- Show that two fields of different characteristic can have neither a product nor a coproduct in the category **Fld** of fields.

Answers: The coproduct of two rings A, B is their tensor product $A \otimes B$, equipped with the ring homomorphisms

$$A \xrightarrow{i_A} A \otimes B : a \mapsto a \otimes 1, \quad B \xrightarrow{i_B} A \otimes B : b \mapsto 1 \otimes b.$$

As a set, $A \otimes B$ is the same thing as the tensor product of A and B as abelian groups; one then gives it a ring structure by defining

$$(a \otimes b)(a' \otimes b') := (aa') \otimes (bb').$$

It is easy to check that the required universal property is satisfied. Perhaps more interesting is to observe that in the more familiar categories \mathbf{Ab} and $R\text{-Mod}$ in which we are used to talking about tensor products, they do *not* arise as colimits, and there is an obvious reason why they shouldn't: the only obviously canonical homomorphisms I can think of from a pair of abelian groups A and B to their tensor product $A \otimes B$ are the trivial ones. The big difference in \mathbf{Ring} is that rings have multiplicative units, and these give rise to canonical *nontrivial* morphisms from A and B to $A \otimes B$ as described above. (For similar reasons, you also should not try to think of tensor products as categorical products—for a more useful categorical perspective on tensor products, see Exercise 1.9.)

The product in \mathbf{Ring} is exactly what you'd expect: the product of rings.

For fields, the problem is that there are in fact no field homomorphisms *at all* between a pair of fields with different characteristics. So for any fields A and B , the need to have morphisms $A, B \rightarrow A \coprod B$ and $A \times B \rightarrow A, B$ means that neither the coproduct nor the product can exist unless A and B have the same characteristic (which their product and coproduct must then also have). For example, \mathbb{Z}_2 and \mathbb{Q} have no coproduct in \mathbf{Fld} , though they do have a coproduct in \mathbf{Ring} , namely $\mathbb{Z}_2 \otimes \mathbb{Q}$, which is an extremely indirect way of writing the trivial ring. (Amusing exercise: show that $1 = 0$ in $\mathbb{Z}_2 \otimes \mathbb{Q}$. The elements 1 and 0 are never equal in a field.)

Exercise 1.5 (initial and terminal objects). In defining limits and colimits of diagrams $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$, the set of objects in \mathcal{J} is not required to be nonempty. When it is empty, we can think of $\text{colim}(\mathcal{F})$ is a coproduct of an empty collection of objects in \mathcal{C} , and $\text{colim}(\mathcal{F})$ is then called an **initial object** in \mathcal{C} . Similarly, the product $\text{lim}(\mathcal{F})$ of an empty collection of objects is called a **terminal** (or **final**) object in \mathcal{C} .

- Reformulate the definitions given above for the terms “initial object” and “terminal object” in a way that makes no reference to limits or colimits, and using this reformulation, give a short proof that both are unique up to canonical isomorphisms, if they exist.
- Show that for any initial object $0 \in \mathcal{C}$, the coproducts $0 \coprod X$ and $X \coprod 0$ exist and the canonical morphisms of X to each are isomorphisms. Similarly, for any terminal object $1 \in \mathcal{C}$, the products $1 \times X$ and $X \times 1$ exist and their canonical morphisms to X are isomorphisms.
- Describe what initial and terminal objects are in each of the following categories, if they exist: \mathbf{Top} , \mathbf{Top}_* , \mathbf{Ab} , \mathbf{Ring} , and \mathbf{Fld} .

Hint: You might guess the last two from Exercise 1.4.

Answers: If \mathcal{J} is the empty category, then there is a unique diagram $\mathcal{F} : \mathcal{J} \rightarrow \mathcal{C}$, but it carries no information. If we want to define a colimit of this diagram, then any object $X \in \mathcal{C}$ can be considered a target; there is no need to specify any morphisms since \mathcal{J} has no objects. The condition of X being a *universal* target is, however, nontrivial: it means that for any other target Y , there is a unique morphism $u : X \rightarrow Y$ such that... well, at this point we would normally say that certain morphisms admit factorizations through the morphism u , but since \mathcal{J} has no objects, there are no morphisms to be factored and thus no further conditions to impose. We are left only with this: $X \in \mathcal{C}$ is an initial object if and only if for every object $Y \in \mathcal{C}$, there is a

unique morphism $X \rightarrow Y$. That's the usual definition—we stated it in a much more roundabout way by talking about coproducts over the empty category.

Here's the dual version: $X \in \mathcal{C}$ is a *terminal object* if and only if for every object $Y \in \mathcal{C}$, there is a unique morphism $Y \rightarrow X$.

With these definitions understood: if $0, 0' \in \mathcal{C}$ are two initial objects, then there is a unique morphism $0 \rightarrow 0'$, and there is also a unique morphism $0' \rightarrow 0$. Moreover, there are unique morphisms $0 \rightarrow 0$ and $0' \rightarrow 0'$, and both of those have to be identity morphisms, since identity morphisms must always exist. It follows that the unique morphisms $0 \rightarrow 0'$ and $0' \rightarrow 0$ are inverse to each other, and are thus isomorphisms. The uniqueness of terminal objects up to unique isomorphisms is proved similarly; there is only a slightly different reason for the uniqueness of the morphisms $1 \rightarrow 1'$ and so forth.

Let's consider the coproduct of an initial object $0 \in \mathcal{C}$ with an arbitrary $X \in \mathcal{C}$. We claim that X itself plays the role of the coproduct, together with the two morphisms

$$\begin{array}{ccc} 0 & & \\ & \searrow & \\ & & X \\ & \nearrow \text{Id} & \\ X & & \end{array}$$

the first of which is determined by the condition that 0 is an initial object. Indeed, suppose Y is given, along with a morphism $f : X \rightarrow Y$ and the unique morphism $0 \rightarrow Y$ (for which there is no freedom of choice). The dashed arrow in the following diagram is then uniquely determined,

$$\begin{array}{ccccc} 0 & & & & \\ & \searrow & & \searrow & \\ & & X & \dashrightarrow & Y \\ & \nearrow \text{Id} & & \nearrow f & \\ X & & & & \end{array}$$

and this establishes the universal property of the coproduct. In this way of representing $0 \coprod X$, the canonical morphism $X \rightarrow 0 \coprod X$ is simply the identity morphism $X \rightarrow X$, and thus an isomorphism. Similar arguments prove the analogous statements about $X \coprod 0$, $1 \times X$ and $X \times 1$.

Here is an inventory of initial and terminal objects in specific categories:

- **Top**: the empty set \emptyset is initial, and every one-point space $*$ is terminal. Note that the initial object in this case is not just unique *up to isomorphism*, but is actually unique, i.e. there really is only one object in **Top** called \emptyset . By contrast, the unique point in a one-point space can be anything, and the collection of all possible one-point spaces is therefore too large to qualify as a set; it is a proper class. Nonetheless, there is indeed a unique homeomorphism between any two of them.
- **Top_{*}**: every one-point space is both an initial and a terminal object.
- **Ab**: every trivial group is both initial and terminal. The answer in **R-Mod** is the same, in case you'd wondered.
- **Ring**: this one's more interesting. According to Exercise 1.4, tensor products are coproducts in **Ring**, so an initial object $R \in \mathbf{Ring}$ should be a ring with the property that $R \otimes A \cong A \cong A \otimes R$ for all rings $A \in \mathbf{Ring}$; plugging in $A := \mathbb{Z}$ as a special case, one deduces $R \cong \mathbb{Z}$. And indeed, for any other ring B , a ring homomorphism $\mathbb{Z} \rightarrow B$ is uniquely determined

by the condition that it preserve the 0 and 1 elements. Terminal objects are trivial rings, i.e. those in which $1 = 0$.

- **Fld**: there are no initial or terminal objects in **Fld**, because as discussed in the answer to Exercise 1.4(c), there do not exist any fields that admit homomorphisms either to or from every other field (of arbitrary characteristic).

Exercise 1.6 (biproducts). Assume \mathcal{A} is a category in which the sets $\text{Hom}(A, B)$ of morphisms $A \rightarrow B$ for each $A, B \in \mathcal{A}$ are equipped with the structure of abelian groups such that composition $\text{Hom}(A, B) \times \text{Hom}(B, C) : (f, g) \mapsto g \circ f$ is always a bilinear map. (Popular examples are the categories **Ab** of abelian groups and **$R\text{-Mod}$** of modules over a commutative ring R .) A **biproduct** of two objects $A, B \in \mathcal{A}$ is an object $C \in \mathcal{A}$ equipped with four morphisms

$$(1.1) \quad \begin{array}{ccc} A & & A \\ & \searrow i_A & \nearrow \pi_A \\ & C & \\ & \nearrow i_B & \searrow \pi_B \\ B & & B \end{array}$$

that satisfy the five relations

$$(1.2) \quad \pi_A i_A = \mathbb{1}_A, \quad \pi_B i_B = \mathbb{1}_B, \quad \pi_A i_B = 0, \quad \pi_B i_A = 0, \quad i_A \pi_A + i_B \pi_B = \mathbb{1}_C.$$

In the categories **Ab** or **$R\text{-Mod}$** , an example of a biproduct of A and B is the direct sum $A \oplus B$ with its canonical inclusion and projection maps. The category \mathcal{A} is called **additive** if every pair of objects has a biproduct.

- Show that for any biproduct as in the diagram (1.1), C with the morphisms i_A, i_B is a coproduct of A and B , and with the morphisms π_A, π_B it is also a product of A and B .
- Show that in the categories **Ab** and **$R\text{-Mod}$** , every biproduct of two objects A, B admits an isomorphism to $A \oplus B$ that identifies the four maps in (1.1) with the obvious inclusions and projections.
- A (covariant or contravariant) functor $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$ between two additive categories is called an **additive functor** if the map defined by \mathcal{F} from $\text{Hom}(A, B)$ to $\text{Hom}(\mathcal{F}(A), \mathcal{F}(B))$ or (in the contravariant case) $\text{Hom}(\mathcal{F}(B), \mathcal{F}(A))$ is a group homomorphism for all $A, B \in \mathcal{A}$. Show that additive functors send all biproducts in \mathcal{A} to biproducts in \mathcal{B} .

Remark: Popular examples of additive functors $\text{Ab} \rightarrow \text{Ab}$ or $R\text{-Mod} \rightarrow R\text{-Mod}$ are $\otimes G$, $G \otimes$, $\text{Hom}(\cdot, G)$ and $\text{Hom}(G, \cdot)$ for any fixed module G , as these arise in the universal coefficient theorems for homology and cohomology.

Answers: Let's show first that (1.1) and (1.2) make C with the morphisms $i_A : A \rightarrow C$ and $i_B : B \rightarrow C$ into a coproduct of A and B . We need to show that the dashed morphism u in the diagram

$$\begin{array}{ccc} A & & X \\ & \searrow i_A & \nearrow f_A \\ & C & \xrightarrow{u} X \\ & \nearrow i_B & \searrow f_B \\ B & & X \end{array}$$

exists and is unique for any given object $X \in \mathcal{A}$ with morphisms f_A, f_B from A and B respectively. Start with uniqueness: if u is a morphism for which this diagram commutes, then using (1.2) and

the assumption that composition is bilinear, we have

$$u = u(i_A\pi_A + i_B\pi_B) = (ui_A)\pi_A + (ui_B)\pi_B = f_A\pi_A + f_B\pi_B.$$

For existence, we then just need to define u by this formula and show that it satisfies $ui_A = f_A$ and $ui_B = f_B$, which also follows easily from the relations (1.2). The proof that C with the morphisms π_A, π_B is a product of A and B is similar.

For part (b), we already know that $A \oplus B$ defines a biproduct of R -modules A and B , so what we really need is a general result about uniqueness of biproducts up to isomorphism. We already have such results for products and coproducts separately, but we cannot directly apply them here, even though we know that biproducts are both; the trouble is that doing so will produce *two* isomorphisms between any two biproducts of A and B , one that arises by viewing them as products, and another by viewing them as coproducts. We want to see that those two isomorphisms are *the same one*.

Concretely, let's suppose that (1.1) and (1.2) are given, and that we also have a second object C' and set of morphisms $i'_A, i'_B, \pi'_A, \pi'_B$ satisfying the same set of relations. We do not need to assume \mathcal{A} is \mathbf{Ab} or $R\text{-Mod}$ for this discussion, as it will make sense in any category for which biproducts can be defined, but some intuition about direct sums may nonetheless be helpful for writing down suitable morphisms between C and C' . Explicitly, define

$$f := i'_A\pi_A + i'_B\pi_B : C \rightarrow C', \quad \text{and} \quad g := i_A\pi'_A + i_B\pi'_B : C' \rightarrow C.$$

Using (1.2), we then have

$$\begin{aligned} gf &= (i_A\pi'_A + i_B\pi'_B)(i'_A\pi_A + i'_B\pi_B) = i_A(\pi'_A i'_A)\pi_A + i_A(\pi'_A i'_B)\pi_B + i_B(\pi'_B i'_A)\pi_A + i_B(\pi'_B i'_B)\pi_B \\ &= i_A\pi_A + i_B\pi_B = \mathbb{1}_C, \end{aligned}$$

and by a similar calculation, $fg = \mathbb{1}_{C'}$, so f is an isomorphism with $g = f^{-1}$. Using f to identify C with C' now transforms the morphism $i_A : A \rightarrow C$ into

$$fi_A = (i'_A\pi_A + i'_B\pi_B)i_A = i'_A(\pi_A i_A) + i'_B(\pi_B i_A) = i'_A : A \rightarrow C',$$

and it transforms the morphism $\pi_A : C \rightarrow A$ into

$$\pi_A f^{-1} = \pi_A(i_A\pi'_A + i_B\pi'_B) = (\pi_A i_A)\pi'_A + (\pi_A i_B)\pi'_B = \pi'_A : C' \rightarrow A,$$

and by similar calculations,

$$fi_B = i'_B, \quad \pi_B f^{-1} = \pi'_B.$$

One can now appeal to abstract principles (i.e. the universal properties of products and coproducts) to deduce that f is indeed the *only* isomorphism $C \rightarrow C'$ that relates the morphisms i_A, i'_A and so forth in this way.

For a covariant additive functor $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$, it is easy to check that \mathcal{F} sends the four morphisms of (1.1) to morphisms

$$\begin{array}{ccc} \mathcal{F}(A) & & \mathcal{F}(A) \\ & \searrow \mathcal{F}(i_A) \quad \mathcal{F}(\pi_A) \nearrow & \\ & \mathcal{F}(C) & \\ & \nwarrow \mathcal{F}(i_B) \quad \mathcal{F}(\pi_B) \searrow & \\ \mathcal{F}(B) & & \mathcal{F}(B) \end{array}$$

in \mathcal{B} that satisfy the five relations (1.2), making $\mathcal{F}(C)$ a biproduct of $\mathcal{F}(A)$ and $\mathcal{F}(B)$. The amusing detail is what happens if \mathcal{F} is *contravariant*: it still works, but the reversal of arrows means that

some roles need to be switched, e.g. the diagram in \mathcal{B} arising from (1.1) must be written as

$$\begin{array}{ccc}
 \mathcal{F}(A) & & \mathcal{F}(A) \\
 & \searrow \mathcal{F}(\pi_A) \quad \mathcal{F}(i_A) \nearrow & \\
 & \mathcal{F}(C) & \\
 & \nearrow \mathcal{F}(\pi_B) \quad \mathcal{F}(i_B) \searrow & \\
 \mathcal{F}(B) & & \mathcal{F}(B)
 \end{array}
 .$$

With $\mathcal{F}(\pi_A), \mathcal{F}(\pi_B)$ now playing the roles formerly played by i_A, i_B and $\mathcal{F}(i_A), \mathcal{F}(i_B)$ playing the roles of π_A, π_B , one easily checks that the five relations (1.2) are satisfied, so $\mathcal{F}(C)$ is again a biproduct of $\mathcal{F}(A)$ and $\mathcal{F}(B)$, with contravariance having transformed inclusions into projections and vice versa.

Exercise 1.7 (fiber products in Diff). As mentioned in lecture, the category Diff of smooth manifolds is one in which many limits and colimits do not exist. An important example is the fiber product of two smooth maps $f : M \rightarrow Q$ and $g : N \rightarrow Q$, which matches the usual topological fiber product

$$M \times_g N := \{(x, y) \in M \times N \mid f(x) = g(y)\} \subset M \times N$$

if the maps f and g are **transverse** to each other (written $f \pitchfork g$), because the implicit function theorem then gives $M \times_g N$ a natural smooth manifold structure for which the obvious projections to M and N are smooth.⁵ If, on the other hand, f and g are not transverse, then the examples below show that all bets are off.

- (a) Suppose $F : P \rightarrow M$ and $G : P \rightarrow N$ are smooth maps that define a target in Diff for the fiber product diagram defined by f and g ; in other words, the diagram

$$\begin{array}{ccc}
 P & \xrightarrow{F} & M \\
 \downarrow G & & \downarrow f \\
 N & \xrightarrow{g} & Q
 \end{array}$$

commutes and consists entirely of smooth manifolds and smooth maps. Interpret this diagram as defining a smooth map

$$u : P \rightarrow M \times N$$

whose image lies in the *topological* fiber product $M \times_g N \subset M \times N$, and show that if F and G satisfy the universal property for a fiber product in Diff, then u is a continuous bijection of P onto $M \times_g N \subset M \times N$.

- (b) Deduce that if $M \times_g N \subset M \times N$ is a smooth submanifold of $M \times N$, then $M \times_g N$ with its projection maps to M and N does in fact define a fiber product in Diff. (Note that this may sometimes hold even if f and g are not transverse.)
- (c) Consider the example $M = N = Q := \mathbb{R}$ with $f(x) := x^2$ and $g(y) := y^2$, thus

$$M \times_g N = \{(x, y) \in \mathbb{R}^2 \mid x^2 = y^2\}.$$

⁵Transversality is a condition on the derivatives of f and g at all points $x \in M$ and $y \in N$ such that $f(x) = g(y) =: p$; writing the derivatives at these points as linear maps $df(x) : T_x M \rightarrow T_p Q$ and $dg(y) : T_y N \rightarrow T_p Q$ between the appropriate tangent spaces, it means that the subspaces $\text{im } df(x)$ and $\text{im } dg(y)$ span all of $T_p Q$. Choosing suitable local coordinates near each point $(x, y) \in M \times_g N$, one can identify $M \times_g N$ locally with the zero-set of a smooth map whose derivative at (x, y) is surjective if and only if the transversality condition holds, so that the implicit function theorem makes $M \times_g N$ a smooth submanifold of $M \times N$.

You will easily convince yourself that this topological fiber product is not a manifold. Show that the pair of maps f, g does not admit any fiber product in Diff . Note that this is a stronger statement than just the observation that $\{x^2 = y^2\} \subset \mathbb{R}^2$ is not an object of Diff . *Hint: You can use parts (a) and (b) to show that if P is a smooth fiber product, then it contains a special point $p \in P$ such that $P \setminus \{p\}$ is diffeomorphic to $\{x^2 = y^2\} \setminus \{(0, 0)\}$.*

- (d) Here's a weirder example: Let $M = Q := \mathbb{R}$, define $N := *$ as a manifold of one point with $g : N \rightarrow Q = \mathbb{R}$ mapping to 0, and choose $f : M = \mathbb{R} \rightarrow \mathbb{R} = Q$ to be any smooth function with

$$f^{-1}(0) = \{-1, -1/2, -1/3, \dots\} \cup \{0\} \cup \{\dots, 1/3, 1/2, 1\}.$$

(If you have doubts about the existence of such a function, try making minor modifications to the function e^{-1/x^2} , or something similar.) Show that in this case, a fiber product in Diff does exist, but is not homeomorphic to the topological fiber product.

Hint: What can you say about continuous maps from locally path-connected spaces to $f^{-1}(0) \subset \mathbb{R}$?

Answers: For part (a), note first that a fiber product diagram in Diff can always also be interpreted as a fiber product diagram in Top , so applying the universal property of the topological fiber product $M \times_{f \times g} N$ immediately gives us a unique *continuous* map $u : P \rightarrow M \times_{f \times g} N$ such that the diagram

$$\begin{array}{ccc} & & M \\ & \nearrow F & \uparrow \\ P & \xrightarrow{u} & M \times_{f \times g} N \\ & \searrow G & \downarrow \\ & & N \end{array}$$

commutes, where the vertical arrows are the obvious projections. This diagram also gives us an explicit formula for u : its composition with the inclusion $M \times_{f \times g} N \hookrightarrow M \times N$ is just

$$(F, G) : P \rightarrow M \times N,$$

which is a smooth map since F and G are smooth, though we cannot sensibly call it a smooth map to $M \times_{f \times g} N$ unless the latter is known to be a smooth submanifold of $M \times N$.

We want to show that if P with the maps F and G satisfies the universal property for a fiber product in Diff , then the map $u : P \rightarrow M \times_{f \times g} N$ described above is a bijection. Indeed, pick any point $(x, y) \in M \times_{f \times g} N$ and consider the pullback diagram

$$\begin{array}{ccc} * & \xrightarrow{x} & M \\ \downarrow y & & \downarrow f \\ N & \xrightarrow{g} & Q \end{array}$$

where the labels “ x ” and “ y ” on arrows are used to indicate the images of maps from a one-point space labelled $*$. The latter is (trivially) a smooth 0-manifold, and the maps defined on it are (trivially) smooth, so this diagram lives in Diff , and the universal property of the fiber product P

therefore produces a unique map $u : * \rightarrow P$ for which the diagram

$$\begin{array}{ccc} & & M \\ & \nearrow x & \uparrow F \\ * & \xrightarrow{u} & P \\ & \searrow y & \downarrow G \\ & & N \end{array}$$

commutes. The image of $u : * \rightarrow P$ is thus the unique point $p \in P$ satisfying $u(p) = (F(p), G(p)) = (x, y)$.

Part (b) follows almost immediately from what was said above: if $M \times_g N$ is a smooth submanifold of $M \times N$, then the map $u : P \rightarrow M \times_g N$ obtained from any smooth fiber product diagram by applying the universal property in **Top** is automatically also smooth, with the consequence that $M \times_g N$ also satisfies the universal property in **Diff**.

For the example in part (c), $M \times_g N \subset \mathbb{R} \times \mathbb{R} = \mathbb{R}^2$ is the union of the two lines $\{y = x\}$ and $\{y = -x\}$, so it is not globally a manifold, though it becomes a smooth 1-manifold if one deletes the singular point $(0, 0)$. Suppose there exists a smooth manifold P and smooth functions $F, G : P \rightarrow \mathbb{R}$ such that the diagram

$$\begin{array}{ccc} P & \xrightarrow{F} & \mathbb{R} \\ \downarrow G & & \downarrow f \\ \mathbb{R} & \xrightarrow{g} & \mathbb{R} \end{array}$$

defines a fiber product in **Diff**. By part (a), the smooth map $(F, G) : P \rightarrow \mathbb{R}^2$ is then a bijection onto the set $\{y = \pm x\}$, so that there is a unique point $p \in P$ with $F(p) = G(p) = 0$. The manifold P must be path-connected, because any point in $\{y = \pm x\}$ can be joined to $(0, 0)$ by a smooth path lying in one of the smooth submanifolds $\{y = x\}$ or $\{y = -x\}$, and the universal property will then produce a smooth map from this submanifold to P , whose image thus contains a path from any given point to p . Now let $\Sigma := \{y = \pm x\} \setminus \{(0, 0)\} \subset \mathbb{R}^2$, defining a smooth 1-dimensional submanifold of \mathbb{R}^2 , and observe that the restrictions to Σ of the two projections $\mathbb{R}^2 \rightarrow \mathbb{R}$ define a smooth fiber product diagram, and thus (since P satisfies the universal property) give rise to a smooth map $u : \Sigma \rightarrow P$, which is inverse to the bijection $P \setminus \{p\} \rightarrow \Sigma$ defined by (F, G) . This shows that $P \setminus \{p\}$ and Σ are diffeomorphic, thus P is a connected smooth manifold that can be turned into a 1-manifold with four connected components by deleting one point. There is no such manifold, so this is a contradiction.

For the example in part (d), we can identify $M \times N = \mathbb{R} \times *$ with \mathbb{R} and thus identify the topological fiber product with the set

$$M \times_g N = f^{-1}(0) \subset \mathbb{R},$$

carrying the subspace topology it inherits as a subset of \mathbb{R} . It is not a manifold, because the point $0 \in f^{-1}(0)$ does not have any connected neighborhood. However, for any given smooth fiber product diagram

$$\begin{array}{ccc} P & \xrightarrow{F} & \mathbb{R} \\ \downarrow & & \downarrow f \\ * & \xrightarrow{0} & \mathbb{R} \end{array},$$

P is a smooth manifold with a smooth function $F : P \rightarrow \mathbb{R}$ whose image is contained in $f^{-1}(0)$, and there is very little freedom in finding functions F with this property: since P is locally path-connected, F must be locally constant. It follows that F does factor through a smooth manifold

with an obvious smooth bijection onto $f^{-1}(0)$: the manifold in question is $f^{-1}(0)$ itself, but with the *discrete* topology instead of the subspace topology. Conclusion: the fiber product in \mathbf{Diff} for our given pair of maps is given by

$$\begin{array}{ccc} f^{-1}(0) & \hookrightarrow & \mathbb{R} \\ \downarrow & & \downarrow f \\ * & \xrightarrow{0} & \mathbb{R} \end{array}$$

where $f^{-1}(0)$ in the corner is understood to carry the discrete topology and is thus a smooth 0-manifold. Its obvious bijection to the topological fiber product ($f^{-1}(0)$ with the subspace topology) is continuous, but not a homeomorphism.

Exercise 1.8. The following bit of abstract nonsense provides a useful tool for proving that objects are isomorphic in various categories, e.g. one can apply it in \mathbf{hTop} to establish homotopy equivalences, or (as in Exercise 1.9 below) to deduce properties of tensor products from a universal property.

In any category \mathcal{C} , each object $X \in \mathcal{C}$ determines a covariant functor

$$\mathrm{Hom}(X, \cdot) : \mathcal{C} \rightarrow \mathbf{Set},$$

which associates to each object $Y \in \mathcal{C}$ the set $\mathrm{Hom}(X, Y)$ of morphisms and to each morphism $f : Y \rightarrow Z$ in \mathcal{C} the map

$$\mathrm{Hom}(X, Y) \xrightarrow{f_*} \mathrm{Hom}(X, Z) : g \mapsto f \circ g.$$

There is similarly a contravariant functor $\mathrm{Hom}(\cdot, X) : \mathcal{C} \rightarrow \mathbf{Set}$ for which morphisms $f : Y \rightarrow Z$ induce maps

$$\mathrm{Hom}(Z, X) \xrightarrow{f^*} \mathrm{Hom}(Y, X) : g \mapsto g \circ f.$$

- (a) Show that for any two objects $X, Y \in \mathcal{C}$, each morphism $f : X \rightarrow Y$ determines a natural transformation $T_f : \mathrm{Hom}(Y, \cdot) \rightarrow \mathrm{Hom}(X, \cdot)$ associating to each object $Z \in \mathcal{C}$ the set map $f^* : \mathrm{Hom}(Y, Z) \rightarrow \mathrm{Hom}(X, Z)$, and that if f is an isomorphism, then the map f^* is bijective for every $Z \in \mathcal{C}$, i.e. T_f is then a *natural isomorphism*.⁶
- (b) Show conversely that every natural transformation $T : \mathrm{Hom}(Y, \cdot) \rightarrow \mathrm{Hom}(X, \cdot)$ is T_f for a unique morphism $f : X \rightarrow Y$, which is an isomorphism of \mathcal{C} if and only if T_f is a natural isomorphism. It follows that X and Y are isomorphic whenever the sets of morphisms $\mathrm{Hom}(X, Z)$ and $\mathrm{Hom}(Y, Z)$ are in bijective correspondence for every third object Z , in a way that is natural with respect to Z .
- (c) Prove contravariant analogues of parts (a) and (b) involving the functors $\mathrm{Hom}(\cdot, X)$ and $\mathrm{Hom}(\cdot, Y)$.

Solution: The interesting step is part (b), so let's just talk about that. (One could give a quick answer to part (a) more or less by mumbling the word “functor”.) Suppose a natural transformation $T : \mathrm{Hom}(Y, \cdot) \rightarrow \mathrm{Hom}(X, \cdot)$ is given, so for every object $Z \in \mathcal{C}$, T defines a set map

$$T_Z : \mathrm{Hom}(Y, Z) \rightarrow \mathrm{Hom}(X, Z)$$

which is required to fit into certain commutative diagrams as dictated by the word “natural”. In particular, choosing $Z := Y$, we observe that T determines a distinguished morphism $f : X \rightarrow Y$ by

$$f := T_Y(\mathrm{Id}_Y) \in \mathrm{Hom}(X, Y).$$

⁶A **natural isomorphism** $T : \mathcal{F} \rightarrow \mathcal{G}$ between two functors $\mathcal{F}, \mathcal{G} : \mathcal{A} \rightarrow \mathcal{B}$ is a natural transformation such that the morphism $T(\alpha) : \mathcal{F}(\alpha) \rightarrow \mathcal{G}(\alpha)$ in \mathcal{B} associated to each object $\alpha \in \mathcal{A}$ is an isomorphism. It follows that T has an inverse, which is also a natural transformation $T^{-1} : \mathcal{G} \rightarrow \mathcal{F}$.

We claim now that, in fact, $T = T_f$. Indeed, given any $Z \in \mathcal{C}$ and $g \in \text{Hom}(Y, Z)$, naturality implies that the diagram

$$\begin{array}{ccc} \text{Hom}(Y, Y) & \xrightarrow{T_Y} & \text{Hom}(X, Y) \\ \downarrow g_* & & \downarrow g_* \\ \text{Hom}(Y, Z) & \xrightarrow{T_Z} & \text{Hom}(X, Z) \end{array}$$

commutes, hence

$$T_Z(g) = T_Z(g \circ \text{Id}_Y) = (T_Z \circ g_*)(\text{Id}_Y) = (g_* \circ T_Y)(\text{Id}_Y) = g_* f = g \circ f = f^* g = T_f(g).$$

Now that we know all natural transformations arise in this way, and after verifying the formula $T_{f \circ g} = T_g \circ T_f$, it follows easily that the morphism $f : X \rightarrow Y$ has an inverse if and only if the corresponding natural transformation T_f has an inverse.

One way to apply this result in homotopy theory is as follows. Suppose we are given a map $f : X \rightarrow Y$ for which we can verify that for all spaces Z , the induced maps

$$f^* : [Y, Z] \rightarrow [X, Z] : g \mapsto g \circ f$$

are bijective. This means that the natural transformation on Hom-functors corresponding to f is a natural isomorphism, therefore implying that f itself is an isomorphism, i.e. the conclusion in this setting is that f is a homotopy equivalence. The variant in part (c) would imply similarly that if the maps

$$f_* : [Z, X] \rightarrow [Z, Y] : g \mapsto f \circ g$$

are known to be bijective for all spaces Z , then f is a homotopy equivalence.

Exercise 1.9 (tensor products). On the category $R\text{-Mod}$ of modules over a commutative ring R , the tensor product satisfies the following universal property: for any three R -modules A, B, C , the natural map

$$\text{Hom}(A \otimes B, C) \xrightarrow{\alpha} \text{Hom}(A, \text{Hom}(B, C)), \quad \alpha(\Phi)(a)(b) := \Phi(a \otimes b)$$

is a bijection. Indeed,

$$\text{Hom}_2(A, B; C) := \text{Hom}(A, \text{Hom}(B, C))$$

can be interpreted as the set of R -bilinear maps $A \times B \rightarrow C$, so the fact that α is bijective means that every such bilinear map factors through the canonical R -bilinear map $A \times B \rightarrow A \otimes B$ and a uniquely determined R -module homomorphism $A \otimes B \rightarrow C$. In fact, α is not just a bijection; it is also an R -module isomorphism, though we will not make use of this fact in the following. The important observation for now is that α defines a natural isomorphism between the two functors $\text{Hom}(\cdot \otimes \cdot, \cdot)$ and Hom_2 from $R\text{-Mod} \times R\text{-Mod} \times R\text{-Mod}$ to \mathbf{Set} , which are contravariant in the first two variables and covariant in the third.

More generally, suppose \mathcal{C} is any category for which the sets $\text{Hom}(X, Y)$ can be regarded as objects in \mathcal{C} for every $X, Y \in \mathcal{C}$, and suppose $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is a functor such that the functors $\mathcal{C} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathbf{Set}$ defined by $\text{Hom}(\cdot \otimes \cdot, \cdot)$ and $\text{Hom}_2 := \text{Hom}(\cdot, \text{Hom}(\cdot, \cdot))$ are naturally isomorphic, so in particular, for every triple of objects $X, Y, Z \in \mathcal{C}$, there is a bijection of sets

$$\text{Hom}(X \otimes Y, Z) \cong \text{Hom}(X, \text{Hom}(Y, Z))$$

that is natural with respect to all three.

- (a) Prove that there is a natural isomorphism relating any two functors $\otimes, \otimes' : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ that satisfy the condition described above. In other words: tensor products are uniquely determined (up to natural isomorphism) by the universal property.

- (b) Prove that \otimes is associative in the sense that the functors $\mathcal{C} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ defined by $(X, Y, Z) \mapsto X \otimes (Y \otimes Z)$ and $(X, Y, Z) \mapsto (X \otimes Y) \otimes Z$ are naturally isomorphic. Prove it using only the universal property, i.e. do not use any knowledge of how \otimes is actually defined in any specific categories.

Solutions: Both parts are applications of Exercise 1.8, which is the right tool for the job because the universal property of \otimes does not tell us what $X \otimes Y$ is, but instead tells us what other functor $\text{Hom}(X \otimes Y, \cdot)$ is naturally isomorphic to, namely $\text{Hom}_2(X, Y; \cdot) := \text{Hom}(X, \text{Hom}(Y, \cdot))$. If we are given two versions \otimes and \otimes' that both satisfy the universal property, we obtain from this a natural isomorphism

$$\text{Hom}(X \otimes Y, \cdot) \cong \text{Hom}(X \otimes' Y, \cdot)$$

for every pair of objects $X, Y \in \mathcal{C}$, and therefore (via Exercise 1.8) an isomorphism $X \otimes Y \cong X \otimes' Y$.

Associativity follows similarly because one can follow two chains of natural bijections that both end at the same destination: for any spaces X, Y, Z, V we have:

$$\text{Hom}(X \otimes (Y \otimes Z), V) \cong \text{Hom}(X, \text{Hom}(Y \otimes Z, V)) \cong \text{Hom}(X, \text{Hom}(Y, \text{Hom}(Z, V))),$$

and also

$$\text{Hom}((X \otimes Y) \otimes Z, V) \cong \text{Hom}(X \otimes Y, \text{Hom}(Z, V)) \cong \text{Hom}(X, \text{Hom}(Y, \text{Hom}(Z, V))).$$

Exercise 1.10 (tensor products of pairs). Let $\mathbf{Top}^{\text{rel}}$ denote the category of pairs of spaces and maps of pairs. When defining the cross and cup products on relative homology and cohomology, one often sees the product of two pairs defined as

$$(X, A) \times (Y, B) = (X \times Y, A \times Y \cup X \times B).$$

- (a) Why is this definition of \times not actually a *product* (in the sense of category theory) on the category $\mathbf{Top}^{\text{rel}}$? What do categorical products in $\mathbf{Top}^{\text{rel}}$ actually look like?
- (b) In the spirit of Exercise 1.9, I would like to argue that \times as defined above should be interpreted as a *tensor product* on $\mathbf{Top}^{\text{rel}}$. Due to some subtle point-set topological issues that I'd rather not get into until next week, it's best for now to dispense with topologies and work instead in the category $\mathbf{Set}^{\text{rel}}$, whose objects are pairs (X, A) of sets with $A \subset X$, and whose morphisms $(X, A) \rightarrow (Y, B)$ are arbitrary (not necessarily continuous) maps $X \rightarrow Y$ that send A into B . In this setting, how can you regard each of the sets $\text{Hom}((X, A), (Y, B))$ as an object of $\mathbf{Set}^{\text{rel}}$ such that there are natural bijections

$$\text{Hom}((X, A) \times (Y, B), (Z, C)) \cong \text{Hom}((X, A), \text{Hom}((Y, B), (Z, C)))$$

for all choices of pairs?

Answers: Categorical products require projection morphisms, but e.g. the projection map $X \times Y \rightarrow X$ does not generally send $A \times Y \cup X \times B$ into A , and thus does not define a map of pairs $(X, A) \times (Y, B) \rightarrow (X, A)$. For a categorical product on $\mathbf{Top}^{\text{rel}}$, the correct definition would be the obvious one,

$$(X, A) \times (Y, B) := (X \times Y, A \times B).$$

If (X, A) and (Y, B) are objects in $\mathbf{Set}^{\text{rel}}$, then $\text{Hom}((X, A), (Y, B))$ also becomes an object in $\mathbf{Set}^{\text{rel}}$ after singling out the subset

$$\{\phi \in \text{Hom}((X, A), (Y, B)) \mid \phi(X) \subset B\} \subset \text{Hom}((X, A), (Y, B)).$$

It is then straightforward to check that set maps of pairs from (X, A) to $\text{Hom}((Y, B), (Z, C))$ are in natural bijective correspondence with set maps of pairs from $(X, A) \times (Y, B)$ to (Z, C) .

The case of this with $A = B = C = \emptyset$ is often written in a more appealing way by using the notation

$$X^Y := \text{Hom}(Y, X) \quad \text{in } \mathbf{Set},$$

so that $\text{Hom}(X \times Y, Z) \cong \text{Hom}(X, \text{Hom}(Y, Z))$ becomes the so-called **exponential law**

$$Z^{X \times Y} \cong (Z^Y)^X.$$

Note that this is one of the few situations in which the categorical product can also sensibly be called a tensor product; they are not the same thing in $\mathbf{Set}^{\text{rel}}$, but in \mathbf{Set} they are.

The reason we removed topologies from the picture before starting this discussion was that one needs to be very careful about defining the right topology on the set $C(X, Y)$ of continuous maps $X \rightarrow Y$ between two spaces if one wants to have a natural bijection

$$C(X \times Y, Z) \cong C(X, C(Y, Z)).$$

In fact, there is *no* right way to define the topology on $C(X, Y)$ so that this works for *all* spaces; one must first restrict the category of spaces under consideration, and then make slight modifications to the definitions of both $C(X, Y)$ and $X \times Y$ as topological spaces. We will go into a little bit of detail about this when it becomes necessary, as without it, one would miss out on some very clever tools coming from stable homotopy theory.

2. WEEK 2

The lecture on 22.04.2024 was cancelled due to illness, so this week contains only one lecture.

Lecture 3 (25.04.2024): The homotopy category and mapping cylinders.

- The homotopy categories \mathbf{hTop} (without base points) and \mathbf{hTop}_* (with base points)
- Notation for diagrams that commute up to homotopy (see the notational glossary above)
- The **double mapping cylinder** of two maps $f : Z \rightarrow X$ and $g : Z \rightarrow Y$,

$$Z(f, g) := (X \amalg (I \times Z) \amalg Y) / \sim, \quad \text{where } (0, z) \sim f(z) \text{ and } (1, z) \sim g(z) \text{ for all } z \in Z.$$

- Role of $Z(f, g)$ as a weak form of pushout in \mathbf{hTop} (it is called a **homotopy pushout**): the diagram

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ g \downarrow & \sim & \downarrow i_X \\ Y & \xrightarrow{i_Y} & Z(f, g) \end{array}$$

commutes up to an obvious homotopy, though not on the nose (the obvious inclusions i_X and i_Y have disjoint images). Diagrams

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ g \downarrow & \sim_H & \downarrow \varphi \\ Y & \xrightarrow{\psi} & Q \end{array}$$

determine maps $Z(f, g) \xrightarrow{u} Q$, constructed in an obvious way out of φ , ψ and the homotopy $\varphi \circ f \xrightarrow{H} \psi \circ g$, so that the diagram

$$\begin{array}{ccc} X & & \\ i_X \downarrow & \searrow \varphi & \\ Z(f, g) & \xrightarrow{u} & Q \\ i_Y \uparrow & \nearrow \psi & \\ Y & & \end{array}$$

commutes (on the nose, i.e. not just up to homotopy).

- Special cases:

- (1) **Mapping cylinder** of $f : X \rightarrow Y$:

$$Z(f) := Z(\text{Id}_X, f) = (I \times Z) \cup_f Y,$$

where the gluing occurs along $\{1\} \times Z$. Convenient feature: $Z(f)$ deformation retracts to Y , so $i_Y : Y \hookrightarrow Z(f)$ is a homotopy equivalence. We can therefore view *every* map $X \rightarrow Y$ “up to homotopy equivalence” as inclusion of a subspace, namely $i_X : X \hookrightarrow Z(f)$. (This trick was used once at the end of *Topologie II*, cf. the last two pages of [Wen23].)

- (2) **Mapping cone** of $f : X \rightarrow Y$: using the unique map $\epsilon : X \rightarrow *$, we define

$$\text{cone}(f) := Z(\epsilon, f) = CX \cup_f Y,$$

where $CX := (I \times X)/(\{0\} \times X)$ is the usual cone of X .

- (3) **Suspension** (unreduced): Not the most direct way to define it, but the familiar suspension SX of a space X is also the double mapping cylinder of a pair of maps from X to one-point spaces:

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & \sim & \downarrow \\ * & \longrightarrow & SX \end{array}.$$

Here the two maps from $*$ to SX have images at the opposite poles, which are points obtained by collapsing $I \times X$ at $\{0\} \times X$ and $\{1\} \times X$ separately.

- Variant for \mathbf{hTop}_* : If X, Y, Z are pointed spaces and f, g are pointed maps, defining a base point on $Z(f, g)$ requires modifying its definition by

$$Z(f, g) := \left(X \vee \frac{I \times Z}{I \times *} \vee Y \right) / \sim, \quad \text{where } (0, z) \sim f(z) \text{ and } (1, z) \sim g(z) \text{ for all } z \in Z.$$

Note: Quotienting $I \times Z$ is necessary because $I \times Z$ on its own has no natural base point, but whenever Z, Z' are two pointed spaces,

$$\text{pointed homotopies } I \times Z \rightarrow Z' \quad \Leftrightarrow \quad \text{pointed maps } \frac{I \times Z}{I \times *} \rightarrow Z'.$$

Everything discussed above has analogues in which all maps are base-point preserving. The pointed version is sometimes called the **reduced** double mapping cylinder, and one can also derive from it special cases such as the **reduced mapping cone** and **reduced suspension**, which we'll have much more to say about later.

- Why is $Z(f, g)$ not really a pushout in \mathbf{hTop} ?

- (1) Our construction of the map $u : Z(f, g) \rightarrow Q$ uses more information than a diagram in \mathbf{hTop} : it uses the actual maps in the diagram (not just their homotopy classes), plus a choice of homotopy. This doesn't mean it cannot work, but is a hint that we may be cheating.

(2) (The real reason): The diagram

$$\begin{array}{ccc}
 X & \xrightarrow{\varphi} & Q \\
 i_X \downarrow & \sim & \downarrow \\
 Z(f, g) & \xrightarrow{u} & Q \\
 i_Y \uparrow & \sim & \uparrow \\
 Y & \xrightarrow{\psi} & Q
 \end{array}$$

does not always uniquely determine $[u] \in [Z(f, g), Q]$. Example: The mapping cone $\text{cone}(\alpha)$ of a degree 2 map $\alpha : S^1 \rightarrow S^1$, say $\alpha(e^{i\theta}) := e^{2i\theta}$ if we think of S^1 as the unit circle in \mathbb{C} . Now $\text{cone}(\alpha) \cong \mathbb{RP}^2$ and the natural inclusion $S^1 \hookrightarrow \text{cone}(\alpha)$ defines the nontrivial element of $\pi_1(\mathbb{RP}^2) = \mathbb{Z}_2$. A homotopy pushout diagram

$$\begin{array}{ccc}
 S^1 & \xrightarrow{\quad} & * \\
 \alpha \downarrow & \sim & \downarrow \\
 S^1 & \xrightarrow{\beta} & Q
 \end{array}$$

now means a choice of space Q and homotopy class $\beta \in [S^1, Q]$ such that $\beta \cdot \beta$ is homotopic to a constant loop. The latter always holds if Q is simply connected, so take $Q := S^2$, and then observe that the diagram

$$\begin{array}{ccc}
 * & \xrightarrow{\quad} & S^2 \\
 \downarrow & \sim & \downarrow \\
 \text{cone}(\alpha) & \xrightarrow{u} & S^2 \\
 \uparrow & \sim & \uparrow \\
 S^1 & \xrightarrow{\quad} & S^2
 \end{array}$$

always commutes up to homotopy, since $[S^1, S^2] \cong * \cong [*, S^2]$. But $[\mathbb{RP}^2, S^2]$ has more than one element, because there exist maps $\mathbb{RP}^2 \rightarrow S^2$ having either possible value of the mod-2 mapping degree (cf. Exercise 2.1).

- Theorem: There exists a category \mathcal{P} whose objects are pushout diagrams (in \mathbf{Top})

$$\begin{array}{ccc}
 Z & \xrightarrow{f} & X \\
 \downarrow g & & \\
 Y & &
 \end{array}$$

such that

- (1) Changing the maps f and g by homotopies produces isomorphic objects of \mathcal{P} ;
- (2) There is a functor $\mathcal{P} \rightarrow \mathbf{hTop}$ sending each pushout diagram to its mapping cylinder $Z(f, g)$.

- Proof sketch: Morphisms in \mathcal{P} are diagrams

$$\begin{array}{ccccc}
 Z & \xrightarrow{f} & X & & \\
 \downarrow g & \searrow \gamma & \sim \phi & \searrow \alpha & \\
 & & Z' & \xrightarrow{f'} & X' \\
 & \searrow \psi & \downarrow g' & & \\
 Y & & & & \\
 & \searrow \beta & & & \\
 & & Y' & &
 \end{array} ,$$

including choices of homotopies ϕ and ψ as part of the data. The notion of composition of such morphisms arises naturally by composing maps and concatenating homotopies.⁷ Such a morphism determines a homotopy pushout diagram

$$\begin{array}{ccc}
 Z & \xrightarrow{f} & X \\
 \downarrow g & \sim H & \downarrow i_{X'} \circ \alpha \\
 Y & \xrightarrow{i_{Y'} \circ \beta} & Z(f', g')
 \end{array}$$

and therefore also an induced map $Z(f, g) \xrightarrow{u} Z(f', g')$. It is a bit tedious but straightforward to check:

- (1) The map induced by a composition of two morphisms in \mathcal{P} is homotopic to the composition of the two induced maps.
- (2) If the maps α, β, γ all have homotopy inverses, one can use them to construct an inverse morphism in \mathcal{P} .

Both only require the same ideas that are needed for proving e.g. that multiplication in the fundamental group is associative. The second point implies, in particular, that the map $Z(f, g) \rightarrow Z(f', g')$ is a homotopy equivalence whenever α, β, γ are.

- Corollary: If $f \sim_h f'$ and $g \sim_h g'$, then $Z(f, g)$ and $Z(f', g')$ are homotopy equivalent.
- Theorem: Pushouts in \mathbf{hTop} and \mathbf{hTop}_* do not always exist.⁸
- Proof sketch in \mathbf{hTop}_* : Fix the obvious base point in S^1 so that our previous degree 2 map $\alpha : S^1 \rightarrow S^1$ preserves base points. A pushout diagram in \mathbf{hTop}_* of the form

$$\begin{array}{ccc}
 S^1 & \longrightarrow & * \\
 \alpha \downarrow & \sim & \downarrow \\
 S^1 & \xrightarrow{\beta} & P
 \end{array}$$

then means a pointed space P together with an element in the 2-torsion subgroup of its fundamental group

$$\beta \in \pi_1(P)_{(2)} := \{\gamma \in \pi_1(P) \mid \gamma^2 = 0\}.$$

⁷It seems likely that I'm oversimplifying this and ought to talk about "homotopy classes of homotopies" if I really want the composition in \mathcal{P} to be associative, but I do not want to give these details more attention than they deserve. I am attempting to present a slightly more highbrow perspective on a sequence of lemmas in [tD08, §4.1–4.2] that seem rather technical and tedious.

⁸...which is why we need to use *homotopy* pushouts instead.

Then P and β satisfy the universal property for a pushout in \mathbf{hTop}_* if and only if for every space Q and $\gamma \in \pi_1(Q)_{(2)}$, the map

$$[P, Q] \rightarrow \pi_1(Q)_{(2)} : u \mapsto u_*\beta$$

is a bijection. Assume this is true, and then consider the surjective map

$$\mathrm{SO}(3) \xrightarrow{p} S^2 : A \mapsto Ae_1,$$

where S^2 is the unit sphere in \mathbb{R}^3 and $e_1, e_2, e_3 \in \mathbb{R}^3$ denotes the standard basis. Taking e_1 as a base point in S^2 , we have

$$p^{-1}(e_1) \cong \mathrm{SO}(2) \cong S^1,$$

giving rise to an exact sequence of pointed spaces

$$S^1 \xrightarrow{i} \mathrm{SO}(3) \xrightarrow{p} S^2.$$

We will see next week that the map $p : \mathrm{SO}(3) \rightarrow S^2$ has a special property: it is a *fibration*, with the consequence that for every space P , the induced sequence of pointed sets

$$[P, S^1] \xrightarrow{i_*} [P, \mathrm{SO}(3)] \xrightarrow{p_*} [P, S^2]$$

is also exact, meaning the preimage of the base point under p_* matches the image of i_* . (Here $[X, Y]$ means the set of homotopy classes of pointed maps $X \rightarrow Y$, so it is a set with an obvious base point.) Combining this with the bijection that we deduced above from the universal property of the pushout, we obtain an exact sequence

$$\pi_1(S^1)_{(2)} \rightarrow \pi_1(\mathrm{SO}(3))_{(2)} \rightarrow \pi_1(S^2)_{(2)},$$

in which the first and last terms both vanish. But $\mathrm{SO}(3) \cong \mathbb{RP}^3$ and thus $\pi_1(\mathrm{SO}(3)) \cong \mathbb{Z}_2$, so the middle term does not vanish, and this is a contradiction.

- To do next week: Define what a fibration is and explain why the sequence of sets of homotopy classes in that proof was exact.

Suggested reading. A more comprehensive treatment of mapping cylinders (including details that I left out of the proof of the theorem about the functor $\mathcal{P} \rightarrow \mathbf{hTop}$) can be found in [tD08, §4.1–4.2]. This does not include the proof that pushouts in \mathbf{hTop}_* don’t exist; I found that in the materials for a course on homotopy theory by Tyrone Cutler, available at <https://www.math.uni-bielefeld.de/~tcutler/> (see the first set of exercises on homotopy pushouts).

Exercises (for the Übung on 2.05.2024).

Exercise 2.1. Review the notions of the \mathbb{Z}_2 -valued and \mathbb{Z} -valued mapping degrees for maps between closed and connected topological manifolds of the same dimension, as covered e.g. in [Wen23, Lecture 35]. Then:

- Show that for every closed and connected topological manifold M of dimension $n \in \mathbb{N}$, the set $[M, S^n]$ contains at least two elements, and infinitely many if M is orientable.
- Does the set $[S^n, M]$ also always have more than one element?

Exercise 2.2. Deduce from the properties of double mapping cylinders the standard fact that there is a functor $S : \mathbf{Top} \rightarrow \mathbf{Top}$ assigning to every space $X \in \mathbf{Top}$ its (unreduced) suspension SX . *Note: This is just intended as a sanity check. There is nothing especially nontrivial to be done here, and there are also more direct ways to show that suspensions define a functor.*

Exercise 2.3. Show that the mapping cone $\mathrm{cone}(f)$ of any homotopy equivalence $f : X \rightarrow Y$ is a contractible space.

Hint: Find a useful morphism in the category \mathcal{P} of pushout diagrams.

Exercise 2.4. Show that for any two maps $f : Z \rightarrow X$ and $g : Z \rightarrow Y$, the singular homologies (with arbitrary coefficients) of the spaces X, Y, Z and $Z(f, g)$ are related by a long exact sequence of the form

$$\dots \rightarrow H_{n+1}(Z(f, g)) \rightarrow H_n(Z) \rightarrow H_n(X) \oplus H_n(Y) \rightarrow H_n(Z(f, g)) \rightarrow H_{n-1}(Z) \rightarrow \dots,$$

and describe explicitly what the two homomorphisms in the middle of this sequence look like. Show that it also works with all homology groups replaced by their reduced counterparts, then write down the special case of a mapping cone and check that what you have is consistent with Exercise 2.3.

Hint: There is a relatively straightforward way to apply the Mayer-Vietoris sequence here, but you could also deduce this as a special case of the exact sequence of the generalized mapping torus derived in [Wen23, Lecture 34].

Exercise 2.5. Prove that pushouts in \mathbf{hTop} do not always exist.

Hint: The proof carried out in lecture for \mathbf{hTop}_ requires only minor modifications. Note that even if X and Y are spaces without base points, the set of homotopy classes $[X, Y]$ still has a natural base point whenever Y is path-connected. (Why?)*

Exercise 2.6. Give explicit examples of homotopic maps

$$f \underset{h}{\sim} f' : Z \rightarrow X \quad \text{and} \quad g \underset{h}{\sim} g' : Z \rightarrow Y$$

such that the mapping cylinders $Z(f, g)$ and $Z(f', g')$ are not homeomorphic. (They will of course be homotopy equivalent!)

Exercise 2.7. The **join** $X * Y$ of two spaces X and Y is the double mapping cylinder $Z(\pi_X, \pi_Y)$ defined via the projection maps $\pi_X : X \times Y \rightarrow X$ and $\pi_Y : X \times Y \rightarrow Y$. Prove that the join of two spheres is always homeomorphic to a sphere: concretely, for every $m, n \in \mathbb{N}$,

$$S^m * S^n \cong S^{m+n+1}.$$

*Hint: Split the double mapping cylinder in half so that you see $S^m * S^n$ as the union of two pieces glued along boundaries that both look like $S^m \times S^n$. Can you think of two compact manifolds that both have $S^m \times S^n$ as boundary? Stare closely at the two pieces, you might recognize them! Now glue them together and ask: what is $S^m * S^n$ the boundary of?*

Exercise 2.8. Many constructions in homotopy theory have analogues in homological algebra, and one of these is the mapping cone. For two chain complexes (A_*, ∂_A) and (B_*, ∂_B) with a chain map $f : A_* \rightarrow B_*$, the **mapping cone of f** is the chain complex $(\text{cone}(f)_*, \partial)$ with

$$\text{cone}(f)_n := A_{n-1} \oplus B_n \quad \text{and} \quad \partial := \begin{pmatrix} -\partial_A & 0 \\ -f & \partial_B \end{pmatrix}.$$

The analogy to the mapping cone in \mathbf{Top} goes through cellular homology: if X, Y are two CW-complexes and $f : X \rightarrow Y$ is a cellular map, then the cone of f inherits a natural cell decomposition whose augmented cellular chain complex $\tilde{C}_*^{\text{CW}}(\text{cone}(f))$ is the cone of the chain map $f_* : \tilde{C}_*^{\text{CW}}(X) \rightarrow \tilde{C}_*^{\text{CW}}(Y)$.⁹

Show that the mapping cone $\text{cone}(f)_*$ of a chain map $f : A_* \rightarrow B_*$ similarly plays the role of a *homotopy pushout* in the category \mathbf{Ch} of chain complexes and chain maps, with the role of a one-point space played by the trivial chain complex $0_* \in \mathbf{Ch}$. Specifically:

⁹This was Problem 2(b) on the take-home midterm for last semester's *Topologie II* course, but for Exercise 2.8, you do not need to know about it.

- (a) There is a natural chain map $i_B : B_* \rightarrow \text{cone}(f)_*$ such that the diagram

$$\begin{array}{ccc} A_* & \longrightarrow & 0_* \\ f \downarrow & \sim & \downarrow \\ B_* & \xrightarrow{i_B} & \text{cone}(f)_* \end{array}$$

commutes up to chain homotopy.

- (b) Any homotopy-commutative diagram in Ch of the form

$$\begin{array}{ccc} A_* & \longrightarrow & 0_* \\ f \downarrow & \sim_H & \downarrow \\ B_* & \xrightarrow{\psi} & D_* \end{array}$$

naturally determines a chain map $u : \text{cone}(f)_* \rightarrow D_*$ such that $u \circ i_B$ is chain homotopic to ψ .

- (c) If we were being strict about the analogy via cellular homology, then the trivial complex 0_* in the diagrams above ought to be replaced by $\tilde{C}_*^{\text{CW}}(*)$, the augmented cellular chain complex of a one-point space, which is not trivial: it has nontrivial entries in degrees 0 and -1 , with the boundary operator giving an isomorphism between them. Explain why this discrepancy does not matter, and nothing in the discussion above would change if we used $\tilde{C}_*^{\text{CW}}(*)$ in place of 0_* .

Hint: None of this is hard... the quickest approach may be by guessing.

3. WEEK 3

Lecture 4 (29.04.2024): Introduction to fibrations.

- The set of (free or pointed) homotopy classes $[X, Y]$ as a pointed set (assuming Y is path-connected in the unpointed case)
- What it means for a sequence of three pointed sets to be exact
- Motivational question: Given a map $p : E \rightarrow B$ and the inclusion $i : F := p^{-1}(*) \hookrightarrow E$, what condition makes the sequence

$$[X, F] \xrightarrow{i_*} [X, E] \xrightarrow{p_*} [X, B]$$

exact for all other spaces X ?

- Definition of the homotopy lifting property (free case) and (free, i.e. unpointed) fibrations $p : E \rightarrow B$. (See next lecture for a precise roundup of the crucial definitions.)
- Terminology: the **base** B and **fibers** $E_b := p^{-1}(b) \subset E$ of a fibration $p : E \rightarrow B$
- Example 1: covering spaces (discrete fibers, lifts of homotopies are *unique*, which does not hold for more general fibrations)
- Example 2: fiber bundles (to be studied later in this course): $\{E_b\}_{b \in B}$ is a continuous family of *homeomorphic* spaces (assuming B is path-connected)
 - Example 2a: For M any smooth n -manifold, its **tangent bundle** $TM = \bigcup_{x \in M} T_x M$ is a fiber bundle whose fibers (the **tangent spaces**) $T_x M$ are all homeomorphic to \mathbb{R}^n . (One can cook up examples with more interesting fibers e.g. by equipping each tangent space with an inner product and taking the unit sphere in each—this produces a fiber bundle with fibers homeomorphic to S^{n-1} , a so-called **sphere bundle**.)
 - Example 2b: The map $p : \text{SO}(3) \rightarrow S^2$ that we used in Lecture 3 for showing that pushouts in hTop_* do not always exist. Observation 1: For base point $e_1 \in S^2$, the fiber $F := p^{-1}(e_1)$ is a subgroup isomorphic to $\text{SO}(2)$, thus homeomorphic to S^1 .

Observation 2: That subgroup acts continuously (in fact smoothly), *freely* and *transitively* from the right on every other fiber, implying that all fibers are homeomorphic (in fact diffeomorphic) to S^1 .

- Remark: When we study fiber bundles in earnest, we will prove that they all have the homotopy lifting property, and are thus fibrations. If you are already familiar with *smooth* fiber bundles and connections, then you should believe this easily for the following reason: any choice of connection on $p : E \rightarrow B$ defines parallel transport maps which uniquely determine a lift of any *smooth* homotopy $X \times I \rightarrow B$. (One has to work harder to also get lifts of all *continuous* homotopies... for this, differential geometry is not enough.)
- Theorem (already proved): For any fibration $p : E \rightarrow B$ with B path-connected, and any space X , the induced sequence of free homotopy classes $[X, F] \rightarrow [X, E] \rightarrow [X, B]$ is exact. (Here the map $F \rightarrow E$ is the inclusion of the fiber $F := p^{-1}(b_0) \subset E$ over any chosen point $b_0 \in B$.)¹⁰
- Idea: If we can show that *every* map $f : X \rightarrow Y$ becomes a fibration after replacing X with some space $X' \simeq X$, then we can do this with the inclusion $F \hookrightarrow E$ and thus extend the exact sequence $[X, F] \rightarrow [X, E] \rightarrow [X, B]$ one more term to the left. Then we can do it again, and again, and extend the sequence as far as we want...
- Example (**path space fibrations**): for $(X, x_0) \in \mathbf{Top}_*$, we define
 - the **free path space**: $C(I, X) := \{\text{continuous maps } I \rightarrow X\}$ with the compact-open topology
 - the **based path space**: $PX := P_{x_0}X := \{\gamma \in C(I, X) \mid \gamma(0) = x_0\}$
 - the **based loop space**: $\Omega X := \Omega_{x_0}X := \{\gamma \in PX \mid \gamma(1) = x_0\}$.

Notice: $C(I, X)$ does not depend on a base point, and it has no natural base point of its own. The spaces PX and ΩX do have natural base points defined by constant paths. Define maps $C(X, I) \xrightarrow{p} X$ and $PX \xrightarrow{p} X$ by $p(\gamma) := \gamma(1)$; for the latter, we notice $p^{-1}(x_0) = \Omega X \subset PX$, making

$$\Omega X \hookrightarrow PX \xrightarrow{p} X$$

an exact sequence of pointed spaces.

- Theorem: (1) $C(I, X) \xrightarrow{p} X$ and $PX \xrightarrow{p} X$ are fibrations. (2) The map $C(I, X) \xrightarrow{p} X$ is also a homotopy equivalence. (3) The space PX is contractible.

The following is a digression, subtitled “The revenge of *Topologie I*”:

- Why is $p : C(I, X) \rightarrow X$ continuous? More generally, is the map

$$\text{ev} : C(X, Y) \times X \rightarrow Y : (f, x) \mapsto f(x)$$

continuous for all spaces X and Y ? (One can show that it is always *sequentially* continuous.)

- Counterexample: $\text{ev} : C(\mathbb{Q}, \mathbb{R}) \times \mathbb{Q} \rightarrow \mathbb{R}$ is not continuous for the obvious (subspace) topology on $\mathbb{Q} \subset \mathbb{R}$. Quick proof: If ev is continuous, then for every continuous $f_0 : \mathbb{Q} \rightarrow \mathbb{R}$, every $x_0 \in \mathbb{Q}$ and every neighborhood $\mathcal{U} \subset \mathbb{R}$ of $y_0 := f_0(x_0)$, there are open neighborhoods $\mathcal{O} \subset C(\mathbb{Q}, \mathbb{R})$ and $\mathcal{W} \subset \mathbb{Q}$ such that $(f, x) \in \mathcal{O} \times \mathcal{W}$ implies $f(x) \in \mathcal{U}$. Without loss of generality, the set $\mathcal{O} \subset C(\mathbb{Q}, \mathbb{R})$ has the form

$$\mathcal{O} = \{f \mid f(K_i) \subset \mathcal{V}_i \text{ for all } i = 1, \dots, N\} \subset C(\mathbb{Q}, \mathbb{R})$$

¹⁰In the lecture I somewhat sloppily asserted that this statement was equally valid in the unpointed and pointed cases, but in fact the pointed case involves some subtleties that I brushed under the rug. These gaps got filled in in Lecture 5.

for some finite collection of compact subsets $K_i \subset \mathbb{Q}$ and open subsets $\mathcal{V}_i \subset \mathbb{R}$, $i = 1, \dots, N$. But since compact subsets of \mathbb{Q} *cannot contain any open subsets*, one can then find two irrational numbers $a < b$ such that

$$\mathcal{W}_0 := (a, b) \cap \mathbb{Q} \subset \mathcal{W}$$

is a nonempty open subset of \mathcal{W} disjoint from $K_1 \cup \dots \cup K_N$. Now define a continuous function $f : \mathbb{Q} \rightarrow \mathbb{R}$ that matches f_0 outside of \mathcal{W}_0 but takes a value $f(x) \notin \mathcal{U}$ for some $x \in \mathcal{W}_0$; this is easy since $a, b \notin \mathbb{Q}$. Then $(f, x) \in \mathcal{O} \times \mathcal{W}$ but $f(x) \notin \mathcal{U}$, a contradiction.

- Message: \mathbb{Q} is a terrible topological space. The main problem: It is not locally compact.
- Lemma 1: If X is locally compact and Hausdorff,¹¹ then $\text{ev} : C(X, Y) \times X \rightarrow Y$ is continuous. (For the proof, see Exercise 3.3.)
- The **exponential law**: For two sets X, Y (not necessarily with topologies), let X^Y denote the set of all (not necessarily continuous) maps $Y \rightarrow X$. Then there is a natural bijection

$$Z^{X \times Y} \cong (Z^Y)^X,$$

identifying each map $f : X \times Y \rightarrow Z$ with the map $\hat{f} : X \rightarrow Z^Y$ defined by $\hat{f}(x)(y) := f(x, y)$.

- Lemma 2: For all topological spaces X, Y, Z , if $f : X \times Y \rightarrow Z$ is continuous, then the corresponding map $\hat{f} : X \rightarrow Z^Y$ is a continuous map into $C(Y, Z)$. The converse also holds if Y is locally compact and Hausdorff. (Proof: see Exercise 3.3.)
- Corollary (since I is locally compact and Hausdorff): Homotopies $X \times I \rightarrow Y$ are naturally equivalent to continuous maps on X with values in the path space $C(I, Y)$.¹²

End of *Topologie I* digression.

- Proof of the theorem on path space fibrations: see [DK01, Theorem 6.15], supplemented by the following remark. In this proof, there are several maps and homotopies to be written down, most of which are pretty straightforward, one just needs to think a little about why they are continuous. Thanks to the digression above, the fact that I is locally compact and Hausdorff ensures this.

Lecture 5 (2.05.2024): Replacing maps with fibrations. This lecture began with some minor extensions and clarifications to the main definition from Lecture 4.

- Definition: A map $p : E \rightarrow B$ has the (free) **homotopy lifting property (HLP)** with respect to some class of spaces $\mathcal{C} \subset \mathbf{Top}$ if the lifting problem

$$\begin{array}{ccc} X & \xrightarrow{\tilde{H}_0} & E \\ \downarrow i_0 & \nearrow \tilde{H} & \downarrow p \\ X \times I & \xrightarrow{H} & B \end{array}$$

¹¹Whether the Hausdorff condition here is truly necessary depends on what definition one takes for the term *locally compact*. I typically define locally compact to mean simply that every point has a compact neighborhood, but many authors (such as tom Dieck [tD08]) prefer a stricter definition in which the compact neighborhood can always be assumed arbitrarily small: concretely, for every point $x \in X$, every neighborhood of x contains a neighborhood of x that is compact. The latter is the condition that one really needs for proving $\text{ev} : C(X, Y) \times X \rightarrow Y$ is continuous, but it is equivalent to the simpler definition whenever X is Hausdorff. I have no plans to consider any examples in which X is not Hausdorff.

¹²I'm not certain, but in the lecture I may have stated this wrongly and said homotopies $X \times I \rightarrow Y$ are equivalent to paths in the space $C(X, Y)$, i.e. maps $I \rightarrow C(X, Y)$. The latter is not true in general unless X is also locally compact and Hausdorff.

is solvable for all $X \in \mathcal{C}$, i.e. given a homotopy H and an initial condition \tilde{H}_0 for a lift, the lifted homotopy \tilde{H} exists. Here i_0 denotes the inclusion $X = X \times \{0\} \hookrightarrow X \times I$.

- Notation convention: For a homotopy $H : X \times I \rightarrow Y$, we will often write

$$H_t := H(\cdot, t) : X \rightarrow Y \quad \text{for each } t \in I.$$

- Definition: $p : E \rightarrow B$ is a **free** (Hurewicz) **fibration** if it satisfies the HLP with respect to all spaces $X \in \mathbf{Top}$. The word “free” (or the synonyms “unpointed” or “unbased”) is included in order to distinguish this from the pointed variant below, but will be omitted whenever possible. The word “Hurewicz” will almost always be omitted, but is meant to distinguish this from certain useful weaker conditions, such as:
- Definition: $p : E \rightarrow B$ is a **Serre fibration** if it satisfies the HLP with respect to all CW-complexes X . Note that E and B do not need to be CW-complexes. This condition is often easier to verify, and has some very nice applications to higher homotopy groups (we’ll get there in a few lectures).
- Definition: A pointed map $p : E \rightarrow B$ has the (pointed) **homotopy lifting property** with respect to some class of pointed spaces $\mathcal{C} \subset \mathbf{Top}_*$ if the lifting problem

$$\begin{array}{ccc} (X, *) & \xrightarrow{\tilde{H}_0} & (E, *) \\ \downarrow i_0 & \nearrow \tilde{H} & \downarrow p \\ (X \times I, \{*\} \times I) & \xrightarrow{H} & (B, *) \end{array}$$

is solvable for all $X \in \mathcal{C}$; in other words, we require the HLP but with maps and homotopies replaced by *pointed maps* and *pointed homotopies*.

- Definition: A pointed map $p : E \rightarrow B$ is a **pointed** (Hurewicz) **fibration** if it satisfies the pointed HLP with respect to all $X \in \mathbf{Top}_*$.
- Theorem (“the main property of fibrations”): Assume $p : E \rightarrow B$ satisfies the (free or pointed) HLP with respect to some class \mathcal{C} in \mathbf{Top} or \mathbf{Top}_* respectively; in the free case, assume also that B is path-connected, so that sets of (free or pointed) homotopy classes $[X, B]$ have natural base points in either case. Denote the inclusion $i : F := p^{-1}(b_0) \hookrightarrow E$, where $b_0 \in B$ is the base point in the pointed case, or any chosen point in the free case. Then for every $X \in \mathcal{C}$, the sequence

$$[X, F] \xrightarrow{i_*} [X, E] \xrightarrow{p_*} [X, B]$$

is exact.

- Convenient fact (see Exercise 3.2): Pointed fibrations are also free fibrations after forgetting their base points.
- Inconvenient fact: If $p : E \rightarrow B$ is a free fibration, choosing base points $* \in B$ and $* \in p^{-1}(*) \subset E$ to make p into a pointed map does *not* automatically make it into a pointed fibration! On the other hand, actual counterexamples are not easy to find, mainly because...
- Sufficiently convenient fact: The aforementioned pointed map $p : E \rightarrow B$ does however satisfy the pointed HLP with respect to all “reasonable” pointed spaces. This means that in practice, one rarely actually needs to worry about the distinction between free and pointed fibrations. (Giving more details on this will require some discussion of cofibrations, which is coming next week.)
- Definition: A sequence of maps $Z \xrightarrow{j} X \xrightarrow{f} Y$ has the **homotopy type of a fibration** if there exists a fibration $p : E \rightarrow B$ with fiber inclusion $i : F := p^{-1}(*) \hookrightarrow E$ and a

homotopy commutative diagram

$$\begin{array}{ccccc} Z & \xrightarrow{j} & X & \xrightarrow{f} & Y \\ \downarrow & \sim & \downarrow & \sim & \downarrow \\ F & \xrightarrow{i} & E & \xrightarrow{p} & B \end{array}$$

in which the vertical maps are all homotopy equivalences. (Note: This definition is sensible in either the free or the pointed case—for the latter, one takes all maps and homotopies to be pointed.) It follows that sequences of the form

$$[Q, Z] \xrightarrow{j^*} [Q, X] \xrightarrow{f^*} [Q, Y]$$

are exact for all Q (assuming as usual in the unpointed case that Y is path-connected). Remark: There are obvious generalizations of this conclusion for cases where $p : E \rightarrow B$ only satisfies the HLP with respect to some smaller class of spaces \mathcal{C} ; then one must also assume $Q \in \mathcal{C}$ in writing down such exact sequences.

- Convention: Unless the words “free” or “pointed” are included explicitly, every statement in the rest of this lecture is meant to be valid for both cases, with closely analogous proofs in either context.
- Theorem 1: For every map $f : X \rightarrow Y$, there exists a space Z (the “**homotopy fiber**” of f) and a map j such that $Z \xrightarrow{j} X \xrightarrow{f} Y$ has the homotopy type of a fibration. In other words, “every map is a fibration up to homotopy equivalence”. Proof at the end of the lecture.
- The dual perspective on the HLP: For topological spaces X, Y , abbreviate

$$Y^X := C(X, Y)$$

with the compact-open topology.¹³ This makes X^I the space of paths in X , and since I is locally compact and Hausdorff, the evaluation map

$$\text{ev} : X^I \times I \rightarrow X : (\gamma, t) \mapsto \gamma(t)$$

is a homotopy between $\text{ev}_0 := \text{ev}(\cdot, 0)$ and $\text{ev}_1 := \text{ev}(\cdot, 1)$; one can deduce from this (see Exercise 3.3) that for every continuous map $f : X \rightarrow Y$, the induced map

$$f^I : X^I \rightarrow Y^I : \gamma \mapsto f \circ \gamma$$

is continuous, thus defining a functor $(\cdot)^I = C(I, \cdot) : \mathbf{Top} \rightarrow \mathbf{Top}$. Moreover, the natural bijection $Y^{X \times I} \cong (Y^I)^X$ identifies homotopies $H : X \times I \rightarrow Y$ with maps $H : X \rightarrow Y^I$ into path space, and this translates the HLP into the diagram

$$\begin{array}{ccccc} X & & \xrightarrow{\quad \widetilde{H}_0 \quad} & & E \\ & \searrow \widetilde{H} & & \searrow \text{ev}_0 & \\ & & E^I & \xrightarrow{\quad \text{ev}_0 \quad} & E \\ & \searrow H & \downarrow p^I & & \downarrow p \\ & & B^I & \xrightarrow{\quad \text{ev}_0 \quad} & B \end{array}$$

Interpretation: the HLP is satisfied if and only if E^I with its maps to E and B^I defines a “*weak fiber product*” of the maps $p : E \rightarrow B$ and $\text{ev}_0 : B^I \rightarrow B$, i.e. the map $X \rightarrow E^I$ is required to exist, but need not be unique (as an actual universal property would require).

¹³We had previously used the notation Y^X to mean *all* (not necessarily continuous) maps $X \rightarrow Y$, but we are now altering the definition of this notation in the context of topological spaces, because it’s a convenient shorthand.

- Constructions of fibrations (proofs are straightforward and mostly consist of drawing some diagrams and adding some dotted arrows):
 - (1) **Projection** maps $B \times F \rightarrow B$ are always fibrations. (Note that here one can clearly see the non-uniqueness of the lifted homotopy, outside of special cases such as when F is a discrete space, which would make the projection a covering map.)
 - (2) **Path space**: By a very slight extension of what we proved last time, the map

$$X^I \xrightarrow{(\text{ev}_0, \text{ev}_1)} X \times X$$

is always a fibration.

- (3) **Compositions**: If $p : E \rightarrow B$ and $f : B \rightarrow A$ are fibrations, then so is $f \circ p : E \rightarrow A$. (Remark: If we didn't already know this, we could now deduce from the first three items on this list that the maps $\text{ev}_0, \text{ev}_1 : X^I \rightarrow X$ individually are also fibrations.)
- (4) **Products**: Given two fibrations $p_i : E_i \rightarrow B_i$ for $i = 1, 2$, the product map $p_1 \times p_2 : E_1 \times E_2 \rightarrow B_1 \times B_2$ is also a fibration.
- (5) **Pullbacks**: Assume E' is a fiber product of $p : E \rightarrow B$ and another map $f : B' \rightarrow B$, so we have a diagram

$$\begin{array}{ccc} E' & \xrightarrow{f'} & E \\ \downarrow p' & & \downarrow p \\ B' & \xrightarrow{f} & B \end{array}$$

and E' can be identified with $B' \times_{f \times p} E \subset B' \times E$ so that f' and p' become the obvious projections. For any $b \in B'$, writing $E'_b := (p')^{-1}(b) \subset E'$, it follows that

$$E'_b \xrightarrow{f'} E_{f(b)}$$

is a homeomorphism, thus we think of E' as a union of the same collection of fibers as E , but parametrized over B' instead of B . Proposition: If p is a fibration, then so is p' . (We then call $p' : E' \rightarrow B'$ the **pullback** of $p : E \rightarrow B$ via the map $B' \rightarrow B$, and sometimes emphasize this by writing $f^*E := E'$. It is also often called an **induced fibration**.) Sketch of proof: Given a homotopy $X \times I \rightarrow B'$, composing it with f gives a homotopy to B , which can be lifted to E . The universal property of the pullback determines from this a unique map $X \times I \rightarrow E'$, which turns out to be the lift we need.

- (6) **Path/loop spaces**: Analogously to the free path space functor $\text{Top} \rightarrow \text{Top} : X \mapsto X^I$, the based path and loop spaces define functors $\text{Top}_* \rightarrow \text{Top}_*$ sending X to PX or ΩX . Proposition: For any free fibration $p : E \rightarrow B$, the map $p^I : E^I \rightarrow B^I$ is also a (free) fibration; similarly for any pointed fibration $p : E \rightarrow B$, the maps $Pp : PE \rightarrow PB$ and $\Omega p : \Omega E \rightarrow \Omega B$ are pointed fibrations. Proof in the free case: The correspondence $Y^{X \times I} \cong (Y^I)^X$ translates the HLP for $p^I : E^I \rightarrow B^I$ with respect to a space X into a lifting problem of the form

$$\begin{array}{ccc} X \times I & \longrightarrow & E \\ \downarrow i_0 \times \text{Id} & & \downarrow p \\ X \times I \times I & \longrightarrow & B \end{array}$$

which is solvable because $p : E \rightarrow B$ has the HLP with respect to $X \times I$. The proofs for the based path and loop spaces are Exercise 3.4.

- Proposition: Every pointed fibration $F \xrightarrow{i} E \xrightarrow{p} B$ (we will often write the inclusion of the fiber $F = p^{-1}(*) \subset E$ as part of the data) determines a canonical pointed homotopy class of maps

$$\Omega B \xrightarrow{\delta} F.$$

Part 1 of the proof: The idea is the same as in covering space theory, where each based loop $\gamma : I \rightarrow B$ gets interpreted as a path and then has a (in this case non-unique) lift $\tilde{\gamma} : I \rightarrow E$ that starts at the base point but may end in some other point of $F = p^{-1}(*)$. Since $p : E \rightarrow B$ has the HLP with respect to the space ΩB , we can do this for all loops at once by interpreting $\text{ev} : \Omega B \times I \rightarrow B$ as a homotopy and lifting it:

$$\begin{array}{ccc} \Omega B & \xrightarrow{\text{const}} & E \\ \downarrow & \nearrow \tilde{\text{ev}} & \downarrow p \\ \Omega B \times I & \xrightarrow{\text{ev}} & B \end{array}$$

On ΩB , ev_0 and ev_1 are both constant maps to the base point of B , thus $\delta := \tilde{\text{ev}}_1 : \Omega B \rightarrow E$ takes values in F . (We will need some machinery developed next week in order to show that the homotopy class of δ is independent of the choice of lift.)

- Theorem 2 (just a preview of our goal for next week, with the caveat that the statement may need minor modifications before it is strictly correct): For any pointed fibration $F \xrightarrow{i} E \xrightarrow{p} B$, every triple of consecutive terms in the sequence of pointed (homotopy classes of) maps

$$\dots \longrightarrow \Omega^2 E \xrightarrow{\Omega^2 p} \Omega^2 B \xrightarrow{\delta} \Omega F \xrightarrow{\Omega i} \Omega E \xrightarrow{\Omega p} \Omega B \xrightarrow{\delta} F \xrightarrow{i} E \xrightarrow{p} B$$

has the homotopy type of a pointed fibration. (Note: Implicit in this sequence is the observation that ΩF has an obvious identification with the fiber of $\Omega E \xrightarrow{\Omega p} \Omega B$ over the base point of ΩB , such that $\Omega i : \Omega F \rightarrow \Omega E$ becomes its inclusion.)

- Remark: Theorem 2 gives us long exact sequences of sets of pointed homotopy classes

$$\dots \rightarrow [X, \Omega^2 B] \rightarrow [X, \Omega F] \rightarrow [X, \Omega E] \rightarrow [X, \Omega B] \rightarrow [X, F] \rightarrow [X, E] \rightarrow [X, B],$$

and since pointed fibrations are also free fibrations, the corresponding sequence of sets of *free* homotopy classes is also exact wherever exactness makes sense (i.e. when the relevant space is known to be path-connected).

- Definition: The (double) **mapping path space** of two maps $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ is

$$P(f, g) := X \times_{f \times \text{ev}_0} Z^I \times_{\text{ev}_1 \times g} Y = \{(x, \gamma, y) \in X \times Z^I \times Y \mid \gamma \text{ is a path in } Z \text{ from } f(x) \text{ to } g(y)\}.$$

This construction is “dual” to the double mapping cylinder, in the sense that it fits into all the same diagrams but with the arrows reversed, e.g. with the obvious projections to X and Y , the diagram

$$\begin{array}{ccc} P(f, g) & \xrightarrow{\pi_X} & X \\ \pi_Y \downarrow & \sim & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

commutes up to an obvious homotopy, and any homotopy commutative diagram of the form

$$\begin{array}{ccc} Q & \xrightarrow{\varphi} & X \\ \psi \downarrow & \sim_H & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

naturally determines a map $u : Q \rightarrow P(f, g)$ for which the diagram

$$\begin{array}{ccc} & & X \\ & \nearrow \varphi & \uparrow \pi_X \\ Q & \xrightarrow{u} & P(f, g) \\ & \searrow \psi & \downarrow \pi_Y \\ & & Y \end{array}$$

commutes. In summary: $P(f, g)$ is a **homotopy pullback** of the maps f and g . (As with homotopy pushouts: We are *not* claiming that $P(f, g)$ defines an actual pullback in the category \mathbf{hTop} , which would require the homotopy class of $u : Q \rightarrow P(f, g)$ above to be determined uniquely by the commutativity (up to homotopy) of the diagram. Exercise for those who are so inclined: show that pullbacks in \mathbf{hTop} do not always exist.)

- Remark: If f and g are pointed maps, then there is an obvious choice of base point for $P(f, g)$ that makes everything in the above discussion pointed. In contrast to the case of mapping cylinders, this does not require any modification to the definition of the space $P(f, g)$ itself.
- Proposition: The map

$$(\pi_X, \pi_Y) : P(f, g) \rightarrow X \times Y$$

is a fibration (and by composition, so therefore are the individual projections π_X and π_Y).

Proof: It's a pullback of the path space fibration $Z^I \rightarrow Z \times Z$:

$$\begin{array}{ccc} P(f, g) & \longrightarrow & Z^I \\ (\pi_X, \pi_Y) \downarrow & & \downarrow (\text{ev}_0, \text{ev}_1) \\ X \times Y & \xrightarrow{f \times g} & Z \times Z \end{array},$$

where the map $P(f, g) \rightarrow Z^I$ is $(x, \gamma, y) \mapsto \gamma$.

- Proof of Theorem 1: Define the **mapping path space** of $f : X \rightarrow Y$ as

$$\begin{aligned} P(f) &:= P(f, \text{Id}_Y) = \{(x, \gamma, y) \in X \times Y^I \times Y \mid \gamma(0) = f(x) \text{ and } \gamma(1) = y\} \\ &= \{(x, \gamma) \in X \times Y^I \mid \gamma(0) = f(x)\}. \end{aligned}$$

By contracting every path back to its starting point, we find a deformation retraction of $P(f)$ to an embedded copy of X , i.e. the map $h : X \rightarrow P(f) : x \mapsto (x, \text{const}_{f(x)})$ is a homotopy inverse of the projection $\pi_X : P(f) \rightarrow X : (x, \gamma) \mapsto x$. Moreover, π_X is a fibration, and more importantly, so is the other projection

$$p := \pi_Y : P(f) \rightarrow Y : (x, \gamma) \mapsto \gamma(1),$$

which now fits into the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{h} & P(f) \\ \downarrow f & \swarrow p & \\ Y & & \end{array},$$

in which h is a homotopy equivalence. One can now take the fiber $Z := p^{-1}(*) \subset P(f)$ with inclusion $i : Z \hookrightarrow P(f)$ and define $j := \pi_X \circ i : Z \rightarrow X$, producing the diagram

$$\begin{array}{ccccc} Z & \xrightarrow{j} & X & \xrightarrow{f} & Y \\ \text{Id} \uparrow & & \pi_X \uparrow & \sim & \uparrow \text{Id} \\ Z & \xrightarrow{i} & P(f) & \xrightarrow{p} & Y \end{array}$$

in which the vertical maps are all homotopy equivalences.

- To do list for next week: Clarify in what sense the fibration $P(f) \xrightarrow{p} Y$ and homotopy fiber Z associated to $f : X \rightarrow Y$ are unique, why the homotopy class of $\delta : \Omega B \rightarrow F$ is well defined, where the long exact sequence in Theorem 2 comes from, why free fibrations with added base points are almost as good in practice as pointed fibrations, and along the way, what a *cofibration* is and what this whole story looks like with all the arrows reversed. That will keep us busy enough.

Suggested reading. The main nontrivial things we did this week can be found in [DK01, §6.2, §6.4 and §6.9]. An unfortunate omission in both [DK01] and [tD08] is the pointed variant of the homotopy lifting property, but there’s a fuller discussion of this and the associated subtleties in the “Fibrations IV” section of Cutler’s lecture notes at <https://www.math.uni-bielefeld.de/~tcutler/>.

Exercises (for the Übung on 16.05.2024). Thursday the 9th is a holiday, so we’ll talk about these exercises (and probably some others) in the Übung for the following week.

Exercise 3.1. The following are two examples of maps $p : E \rightarrow B$ with the property that all fibers $E_b := p^{-1}(b)$ are homotopy equivalent—we will see next week that this is a property that fibrations must have, though in these examples, the fibers are *not* all homeomorphic, so they cannot be fiber bundles. Determine whether each is actually a fibration.

- The projection $E \rightarrow \mathbb{R} : (x, y) \mapsto x$ of the subset $E := \{(x, y) \in \mathbb{R}^2 \mid |y| \leq |x|\}$.
- The projection $E \rightarrow I : (x, y) \mapsto x$ of the subset $E := (I \times \{0\}) \cup (\{0\} \times I)$.

Exercise 3.2. Prove that every pointed fibration becomes a free fibration after forgetting the base points.

Hint: For any $X \in \mathbf{Top}$ and $Y \in \mathbf{Top}_*$, unpointed maps $X \rightarrow Y$ are equivalent to pointed maps $X_+ \rightarrow Y$, for a pointed space X_+ defined as the disjoint union of X with a one point space.

Exercise 3.3. For this exercise, let’s agree to call a space X **locally compact** if every neighborhood of every point $x \in X$ contains a compact neighborhood of x .¹⁴ If you prefer the convention that “locally compact” just means every point has a compact neighborhood, then feel free to add the assumption that X is Hausdorff, which makes the simpler definition of locally compact equivalent to the stricter one stated above. We assume as usual that the space $C(X, Y)$ of continuous maps $X \rightarrow Y$ carries the compact-open topology. The first three parts below add up to the proofs of two lemmas that were stated without proof in lecture.

- Prove that if X is locally compact, then the evaluation map $\text{ev} : C(X, Y) \times X \rightarrow Y : (f, x) \mapsto f(x)$ is continuous.

¹⁴This definition presumes the term **neighborhood of x** to mean any set that contains an open set containing x , i.e. the neighborhood itself need not be open.

- (b) Prove that for any spaces X, Y, Z and any continuous map $f : X \times Y \rightarrow Z$, the map $\hat{f} : X \rightarrow C(Y, Z)$ defined by $\hat{f}(x)(y) := f(x, y)$ is also continuous, thus defining an injective map

$$(3.1) \quad C(X \times Y, Z) \rightarrow C(X, C(Y, Z)) : f \mapsto \hat{f}.$$

Remark: One would ideally also like to know that the map (3.1) is continuous, but let's not worry about that for now.

- (c) Prove that for two given spaces Y and Z , the evaluation map $\text{ev} : C(Y, Z) \times Y \rightarrow Z$ is continuous if and only if the map (3.1) is surjective for all spaces X .

Hint: The identity map is continuous on all spaces.

Comment: It follows in particular that (3.1) is a bijection whenever Y is locally compact; we have already made ample use of the special case $Y := I$ in the lectures.

- (d) Give a concrete example of three spaces for which the map (3.1) is not surjective.
 (e) Writing $X^I := C(I, X)$ for the space of paths in X , show that for any continuous map $f : X \rightarrow Y$, the induced map $f^I : X^I \rightarrow Y^I : \gamma \mapsto f \circ \gamma$ is continuous.

Before we continue, here is a definition: A continuous map $q : \tilde{X} \rightarrow X$ is called a **quotient map** if it is surjective and the open sets $\mathcal{U} \subset X$ are precisely the sets for which $q^{-1}(\mathcal{U}) \subset \tilde{X}$ is open. Equivalently, q is a quotient map if and only if it descends to a homeomorphism $\tilde{X}/\sim \rightarrow X$, for the equivalence relation \sim on \tilde{X} such that $x \sim y$ means $q(x) = q(y)$. Most crucially, being a quotient map means that in any diagram of the form

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow q & \searrow \tilde{f} & \\ X & \xrightarrow{f} & Y, \end{array}$$

continuity of the map \tilde{f} implies that f is also continuous. (The converse is of course obvious, since q is continuous.)

- (f) Given two quotient maps $p : \tilde{X} \rightarrow X$ and $q : \tilde{Y} \rightarrow Y$, can you show that the product map $p \times q : \tilde{X} \times \tilde{Y} \rightarrow X \times Y$ is also a quotient map? Give it a try, but do not try too hard. . . Once you've gotten stuck and realized that it isn't obvious, take a look at [Mun75, pp. 143–144].
 (g) Prove that if Y is a space with the property that $\text{ev} : C(Y, Z) \times Y \rightarrow Z$ is continuous for every space Z , then for every quotient map $q : \tilde{X} \rightarrow X$, the product

$$q \times \text{Id}_Y : \tilde{X} \times Y \rightarrow X \times Y$$

is also a quotient map. In particular, this is true whenever Y is locally compact.

- (h) In last week's Übung, I sketched an approach to proving $S^m * S^n \cong S^{m+n+1}$ (Exercise 2.7) that led to the more general formula

$$X * Y \cong (CX \times Y) \cup_{X \times Y} (X \times CY),$$

obtained by splitting the double mapping cylinder in the middle and reinterpreting the quotients that one sees in the two halves. I also mentioned however that it is not so obvious how generally this formula holds, because e.g. $CX \times Y$ is a product of a quotient, which is not always homeomorphic to the corresponding quotient of a product. Can you name some conditions on X and Y that will guarantee that the formula holds? (Your conditions should preferably include the special case with $X = S^m$ and $Y = S^n$!)

Exercise 3.4. In lecture, we exploited the natural bijective correspondence between maps $X \rightarrow Y^I$ and maps $X \times I \rightarrow Y$ to prove that for any fibration $p : E \rightarrow B$, the map $p^I : E^I \rightarrow B^I$ is also a fibration, give or take some minor details (e.g. the continuity of p^I is Exercise 3.3(e) above).

- (a) Describe a pointed space $P'X$ associated to every pointed space X with the property that there is a natural bijective correspondence between pointed maps $X \rightarrow PY$ to the based path space and pointed maps $P'X \rightarrow Y$. Moreover, there should also be a bijective correspondence between pointed homotopies $X \times I \rightarrow PY$ and pointed homotopies $P'X \times I \rightarrow Y$.

Achtung: The detail about homotopies will require you to think about products of quotients, so Exercise 3.3(g) may be useful.

- (b) Do the same thing as in part (a) for pointed maps/homotopies to the based loop space ΩY .
(c) Prove the result stated in lecture that for any pointed fibration $p : E \rightarrow B$, the induced maps $Pp : PE \rightarrow PB$ and $\Omega p : \Omega E \rightarrow \Omega B$ are also pointed fibrations.

Exercise 3.5. Prove that if X is path-connected, then the homotopy type of the based loop space ΩX is independent of the choice of base point.

Exercise 3.6. Formulate an analogue for mapping path spaces $P(f, g)$ of the theorem we previously proved about mapping cylinders $Z(f, g)$ defining a functor from a category of pushout diagrams to \mathbf{hTop} . Convince yourself in this way that the homotopy type of $P(f, g)$ only depends on the homotopy classes of the two maps $f : X \rightarrow Z$ and $g : Y \rightarrow Z$.

Exercise 3.7. The mapping path space $P(f) = \{(x, \gamma) \in X \times Y^I \mid \gamma(0) = f(x)\}$ of a map $f : X \rightarrow Y$ can be described as the fiber product of the maps $f : X \rightarrow Y$ and $\text{ev}_0 : Y^I \rightarrow Y$, so by the universal property of the fiber product, the diagram

$$\begin{array}{ccc} X^I & \xrightarrow{f^I} & Y^I \\ \downarrow \text{ev}_0 & & \downarrow \text{ev}_0 \\ X & \xrightarrow{f} & Y \end{array}$$

determines a map $u : X^I \rightarrow P(f)$. Show that $f : X \rightarrow Y$ is a fibration if and only if the map $u : X^I \rightarrow P(f)$ admits a right-inverse $\lambda : P(f) \rightarrow X^I$; in this situation, λ is sometimes called a **lifting function** for the fibration $f : X \rightarrow Y$.

4. WEEK 4

Thursday this week is a holiday, so there is only one lecture and no Übung.

Lecture 6 (6.05.2024): The transport functor.

- Recall: We constructed for every (unpointed or pointed) map $f : X \rightarrow Y$ a diagram

$$\begin{array}{ccccc} F(f) & \xrightarrow{j} & X & \xrightarrow{f} & Y \\ \text{Id} \uparrow & & \pi_X \uparrow & \sim & \uparrow \text{Id} \\ F(f) & \xrightarrow{i} & P(f) & \xrightarrow{p} & Y \end{array}$$

where the bottom row is a (free or pointed) fibration with fiber $F(f) := p^{-1}(*) \subset P(f)$ (preimage of the base point $* \in Y$ if pointed, an arbitrary point if not), and all the vertical maps are homotopy equivalences. We call $F(f)$ the **homotopy fiber** of $f : X \rightarrow Y$. We also had a homotopy inverse $h : X \rightarrow P(f)$ of π_X fitting into the diagram

$$\begin{array}{ccccc} F & \hookrightarrow & X & \xrightarrow{f} & Y \\ \downarrow h & & \downarrow h & & \downarrow \text{Id} \\ F(f) & \hookrightarrow & P(f) & \xrightarrow{p} & Y \end{array}$$

which commutes on the nose (not just up to homotopy), where $F := f^{-1}(*) \subset X$, thus defining a **comparison map**

$$F \xrightarrow{h} F(f)$$

from the “actual” fiber of f to its homotopy fiber.

- Question: Does $f : X \rightarrow Y$ uniquely determine (up to what notion of equivalence?) the fibration $p : E \rightarrow Y$ in any diagram of the form

$$\begin{array}{ccc} X & \xrightarrow{\sim_{h.e.}} & E \\ & \searrow f \quad \swarrow p & \\ & Y & \end{array} \quad ?$$

For instance, if $f : X \rightarrow Y$ is already a fibration, are the two fibrations (and thus their fibers) equivalent in some sense?¹⁵

- Inspiration from differential geometry: For a smooth fiber bundle $p : E \rightarrow B$, any choice of *connection* associates to each smooth path $x \rightsquigarrow y$ in B a *parallel transport* diffeomorphism

$$E_x \xrightarrow{P_\gamma} E_y,$$

and it is compatible with smooth concatenation of paths: $P_{\alpha\beta} = P_\beta \circ P_\alpha$. Connections live in a contractible space of choices, so up to homotopy, P_γ is independent of this choice and depends only on the (smooth) homotopy class of the path γ . Given any smooth homotopy $H : X \times I \rightarrow B$ between maps $H_0, H_1 : X \rightarrow B$, parallel transport determines a correspondence

$$\begin{array}{ccc} \text{lifts} & \begin{array}{ccc} & \nearrow \tilde{H}_0 & \\ X & \xrightarrow{H_0} & B \\ & \searrow p & \end{array} & \mapsto & \begin{array}{ccc} & \nearrow \tilde{H}_1 & \\ X & \xrightarrow{H_1} & B \\ & \searrow p & \end{array} \end{array}$$

defined by $\tilde{H}_1(x) := P_{H(x,\cdot)} \circ \tilde{H}_0(x)$. At the level of homotopy classes of lifts, this correspondence is independent of choices, and depends on H only up to (smooth) homotopy of homotopies. In homotopy theory, we have no smooth structures and cannot talk about connections... but we probably *can* prove that things are unique up to homotopy!

- Definition: Given $B \in \mathbf{Top}$, the category \mathbf{Top}_B of **spaces over B** has objects that are pairs (X, f) with X a space and $f : X \rightarrow B$ a map, and the set of morphisms $\mathrm{Hom}((X, f), (Y, g))$ consists of **maps over B** , meaning maps $\varphi : X \rightarrow Y$ that fit into the diagram

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & Y \\ & \searrow f \quad \swarrow g & \\ & B & \end{array} \quad .$$

Two such morphisms φ, ψ are **homotopic over B** if there exists a homotopy $\varphi \xrightarrow{H} \psi$ such that H_t is a morphism $(X, f) \rightarrow (Y, g)$ for every $t \in I$. This notion defines the corresponding homotopy category \mathbf{hTop}_B , and isomorphisms in this category are called **homotopy equivalences over B** . There are similar definitions for categories $\mathbf{Top}_{B,*}$ and $\mathbf{hTop}_{B,*}$ in which all maps and homotopies are required to be pointed.

¹⁵In the lecture I stated this question a bit differently, involving a more complicated diagram, but I later realized that that version was not exactly the question we are going to answer, nor is it the one that we really *need* to answer.

- Notation: Given two objects X, Y in \mathbf{Top}_B or $\mathbf{Top}_{B,*}$, we denote by

$$[X, Y]_B := \text{Hom}(X, Y) \text{ in } \mathbf{hTop}_B \text{ or } \mathbf{hTop}_{B,*} \text{ resp.}$$

the set of homotopy classes of (unpointed or pointed) maps $X \rightarrow Y$ over B . We can also write $[(X, f), (Y, g)]_B$ whenever the maps $f : X \rightarrow B$ and $g : Y \rightarrow B$ defining these objects need to be specified.

- Definition: Given $X, Y \in \mathbf{Top}$, the **homotopy groupoid** $\Pi(X, Y)$ is a category whose objects are maps $f : X \rightarrow Y$, with morphisms

$$\text{Hom}(f, g) := \left\{ \text{homotopies } f \xrightarrow{H} g \right\} / \sim,$$

where the equivalence relation is “homotopy of homotopies”: $H \sim H'$ means there is a homotopy $H \xrightarrow{\Phi} H'$ of maps $X \times I \rightarrow Y$ such that $\Phi_s := \Phi(\cdot, \cdot, s) : X \times I \rightarrow Y$ for each $s \in I$ is also a homotopy $f \xrightarrow{\Phi_s} g$. Composition of morphisms is defined by concatenation of homotopies. (Easy exercise: The equivalence relation makes this notion of composition associative. The proof is essentially the same as the proof that multiplication in the fundamental group is associative.)

- Remark: $\Pi(X, Y)$ is called a *groupoid* (and not just a category) because all of its morphisms are invertible; one can always reverse homotopies.
- Special case: $\Pi(Y) := \Pi(*, Y)$ is the **fundamental groupoid** of Y , and for each $y \in Y$, $\text{Hom}(y, y)$ is then the (opposite of the) fundamental group $\pi_1(Y, y)$.¹⁶
- For $X, Y \in \mathbf{Top}_*$, there is a pointed variant of $\Pi(X, Y)$ whose objects are pointed maps and morphisms are homotopy classes of pointed homotopies. Amusing exercise: Is $\Pi(*, Y)$ interesting in the pointed case?
- Theorem: For every (free or pointed) fibration $p : E \rightarrow B$ and every space X (unpointed or pointed), there is a well-defined **transport functor**

$$\Pi(X, B) \rightarrow \mathbf{Set}$$

which associates to each map $f : X \rightarrow B$ the set $[(X, f), (E, p)]_B$ of homotopy classes of maps over B ; we can interpret these as homotopy classes of *lifts* $\tilde{f} : X \rightarrow E$ of $f : X \rightarrow B$. To each homotopy class of homotopies $f \xrightarrow{H} g$ of maps $f, g : X \rightarrow B$, it associates the map

$$[(X, f), (E, p)]_B \xrightarrow{H_\#} [(X, g), (E, p)]_p$$

which sends the homotopy class of the lift \tilde{f} to the homotopy class of a lift \tilde{g} obtained by lifting $H : X \times I \rightarrow B$ to a homotopy $\tilde{H} : X \times I \rightarrow E$ from \tilde{f} to \tilde{g} .

- Remark: It is educational to try using the HLP to prove that $H_\#$ is independent of choices, but you will get stuck at some point and notice that the lifting problem you need to solve is more complicated than the one addressed by the HLP. We will deal with this next week, after talking a bit about the homotopy *extension* property and cofibrations. For the rest of this lecture, we take the existence of the transport functor as a black box and explore some of its applications.

¹⁶A slightly annoying detail here is that while $\text{Hom}(y, y)$ has a natural group structure defined by composition of morphisms—which in this case means homotopy classes of concatenation of paths—the conventions of category theory then force multiplication in $\text{Hom}(y, y)$ to be defined by $[\alpha][\beta] := [\beta \cdot \alpha]$. This is why, strictly speaking $\text{Hom}(y, y)$ is the *opposite* group of $\pi_1(Y, y)$, rather than $\pi_1(Y, y)$ itself. For any group G with multiplication of elements $g, h \in G$ denoted by $gh \in G$, the **opposite group** G^{op} can be defined as the same set but with a new multiplication law “ \cdot ” defined by $g \cdot h := hg$, so there is no difference if G happens to be abelian, but in general G and G^{op} are different (though isomorphic!) groups. One occasionally sees claims in the literature that the “correct” definition of $\pi_1(Y, y)$ really should be what we normally call $\pi_1(Y, y)^{\text{op}}$, so that it matches $\text{Hom}(y, y)$ rather than its opposite group. But this idea does not seem to have caught on.

- Theorem: For any fibration $p : E \rightarrow B$ and any two homotopic maps $f_0, f_1 : B' \rightarrow B$, the pullback fibrations $f_0^*E \rightarrow B'$ and $f_1^*E \rightarrow B'$ are homotopy equivalent over B' . It follows in particular that for every $b \in B'$, there is a homotopy equivalence $(f_0^*E)_b \rightarrow (f_1^*E)_b$ between corresponding fibers.
- Corollary (the case $B' := *$): For a fibration $p : E \rightarrow B$, any two fibers over the same path-component of B are homotopy equivalent.
- Proof of the theorem: For each $i = 0, 1$ we have pullback diagrams

$$\begin{array}{ccc} f_i^*E & \xrightarrow{f'_i} & E \\ \downarrow p_i & & \downarrow p \\ B' & \xrightarrow{f_i} & B \end{array},$$

Let $B' \xrightarrow{f_t} B$ for $t \in I$ denote the family of maps defined by a given homotopy

$$f_0 \xrightarrow{F} f_1.$$

The family $f_0^*E \xrightarrow{f_t \circ p_0} B$ then defines a homotopy

$$f_0 \circ p_0 \xrightarrow{H} f_1 \circ p_0$$

of maps $f_0^*E \rightarrow B$, and using the transport functor, we obtain a bijection $H_\#$ that associates to each homotopy class of lifts of $f_0 \circ p_0$ a homotopy class of lifts of $f_1 \circ p_0$. Since $f'_0 : f_0^*E \rightarrow E$ is a lift of $f_0 \circ p_0 : f_0^*E \rightarrow B$, we can feed this into $H_\#$ and thus obtain a lift $g : f_0^*E \rightarrow E$ of $f_1 \circ p_0$, and by the universal property of the pullback f_1^*E , this uniquely determines the map $\Phi_F : f_0^*E \rightarrow f_1^*E$ in the following diagram

$$\begin{array}{ccccc} f_0^*E & & & & \\ & \searrow \Phi_F & & \nearrow g & \\ & & f_1^*E & \xrightarrow{f'_1} & E \\ & \searrow p_0 & \downarrow p_1 & & \downarrow p \\ & & B' & \xrightarrow{f_1} & B \end{array}.$$

By Exercise 4.1 below, this construction defines a functor

$$\Pi(B', B) \rightarrow \mathbf{hTop}_{B'}$$

which associates to each map $f : B' \rightarrow B$ the induced fibration $f^*E \rightarrow B'$ and to each homotopy class of homotopies $f \xrightarrow{F} g$ the homotopy class of maps over B' represented by $\Phi_F : f^*E \rightarrow g^*E$ as constructed above via the transport functor. Since morphisms in $\Pi(B', B)$ are all invertible, the maps Φ_F obtained in this way are all isomorphisms in $\mathbf{hTop}_{B'}$, meaning homotopy equivalences over B' .

- Theorem: If $E \xrightarrow{p} B$ and $E' \xrightarrow{p'} B$ are two fibrations and $f : E \rightarrow E'$ is a homotopy equivalence of spaces that is also a map over B (with respect to p, p'), then f is also a homotopy equivalence over B .

- Remark: Using Exercise 4.2 below, it follows that whenever we have two ways of replacing a map $f : X \rightarrow Y$ by fibrations $p_i : E_i \rightarrow Y$ as in the diagram

$$\begin{array}{ccccc} E_0 & \xleftarrow{\sim_{h.e.}} & X & \xrightarrow{\sim_{h.e.}} & E_1 \\ & \searrow \sim & \downarrow f & \swarrow \sim & \\ & p_0 & Y & p_1 & \end{array},$$

the two fibrations must be homotopy equivalent over Y , and their corresponding fibers therefore homotopy equivalent. In particular, all reasonable definitions of the term “homotopy fiber” give the same thing up to homotopy equivalence.

- Preparation for the proof: Given a map $X \xrightarrow{f} Y$ and a space E , f induces a map

$$[Y, E] \xrightarrow{f^*} [X, E] : \varphi \mapsto f^* \varphi := \varphi \circ f,$$

which is obviously bijective if f is a homotopy equivalence. If we are also given maps $q : Y \rightarrow B$ and $p : E \rightarrow B$, then for any map $\varphi : Y \rightarrow E$ over B , the diagram

$$\begin{array}{ccccc} & & f^* \varphi := \varphi \circ f & & \\ & \nearrow f & & \searrow \varphi & \\ X & \xrightarrow{\quad} & Y & \xrightarrow{\quad} & E \\ & \searrow f^* q := q \circ f & \downarrow q & \swarrow p & \\ & & B & & \end{array}$$

means that f also induces a map

$$[(Y, q), (E, p)]_B \xrightarrow{f^*} [(X, f^* q), (E, p)]_B.$$

- Lemma: In the situation above, if $p : E \rightarrow B$ is a fibration and $f : X \rightarrow Y$ is a homotopy equivalence, then the map $[(Y, q), (E, p)]_B \xrightarrow{f^*} [(X, f^* q), (E, p)]_B$ is also bijective.
- Proof of the lemma: Given a homotopy inverse $g : Y \rightarrow X$ of X , choose a homotopy $\text{Id}_Y \xrightarrow{H} f \circ g$, so that $q \circ H$ is then a homotopy of maps $Y \rightarrow B$ from q to $q \circ f \circ g = g^* f^* q$. We claim that the diagram

$$(4.1) \quad \begin{array}{ccccc} & & (q \circ H)_\# & & \\ & \nearrow & & \searrow & \\ [(Y, q), (E, p)]_B & \xrightarrow{f^*} & [(X, f^* q), (E, p)]_B & \xrightarrow{g^*} & [(Y, g^* f^* q), (E, p)]_B \end{array}$$

commutes. The reason is that for any given map $(Y, q) \xrightarrow{\varphi} (E, p)$ over B , the following diagram reveals that there is an obvious choice of lift for the homotopy $q \circ H : Y \times I \rightarrow B$ with initial condition $\varphi : Y \rightarrow E$:

$$\begin{array}{ccccc} Y & \xrightarrow{\quad \varphi \quad} & E & & \\ & \searrow \text{Id} & \nearrow \varphi & & \\ & & Y & & \\ & \nearrow H & \searrow q & & \\ Y \times I & \xrightarrow{\quad q \circ H \quad} & B & & \end{array}$$

Choosing $\varphi \circ H : Y \times I \rightarrow E$ as the lifted homotopy, it defines a homotopy from φ to $\varphi \circ f \circ g = g^* f^* \varphi$ and thus proves the claim. Since $(q \circ H)_\#$ is a bijection, it follows that

f^* is injective and g^* is surjective. Using a homotopy of $g \circ f$ to Id_X , one can apply the same trick again to show that the composition

$$[(X, f^*q), (E, p)]_B \xrightarrow{g^*} [(Y, g^*f^*q), (E, p)]_B \xrightarrow{f^*} [(X, f^*g^*f^*q), (E, p)]_B$$

is also bijective, implying that the same map g^* is also injective, and thus bijective. Since the composition g^*f^* in (4.1) is bijective, it now follows that f^* is bijective.

- Remark: The proof of the lemma should remind you of the proof that homotopy equivalences induce isomorphisms of fundamental groups (in spite of the annoying detail that the homotopy inverse need not respect base points). In fact, there is a dual version of this lemma for cofibrations, a special case of which involves homotopy classes of maps $S^1 \rightarrow X$ over a one point space, and the result in that case is precisely the isomorphism of fundamental groups.
- The proof of the theorem about homotopy equivalence of fibrations now follows from abstract nonsense; see Exercise 4.3 below.

Suggested reading. The notions of “spaces/maps over B ” and the homotopy groupoid are introduced in [tD08, §2.2 and §2.9], with the special case of the fundamental groupoid treated at length in §2.5. My presentation of the transport functor is based essentially on [tD08, §5.6], though tom Dieck only gives very brief sketches of proofs in that section, since it appears after the corresponding discussion about cofibrations (which is formally similar).

In [DK01, §6.6], you will also find a fairly down-to-earth proof of the fact that for the fibration $P(f) \rightarrow Y$ constructed out of the mapping path space of any map $f : X \rightarrow Y$, the associated homotopy equivalence $h : X \rightarrow P(f)$ is also a homotopy equivalence over Y whenever $f : X \rightarrow Y$ itself is a fibration. This is less general than what we proved, because it applies only to a specific fibration $P(f) \rightarrow Y$ rather than an arbitrary fibration over Y that fits into a suitable diagram with $f : X \rightarrow Y$. Unfortunately, the proof of the main theorem about the long exact fibration sequence in [DK01, §6.11] sneakily uses the more general version of this uniqueness result, so as far as I can tell, this is a logical gap in the book.

Exercises (also for the Übung on 16.05.2024).

Exercise 4.1. In lecture we used the transport functor to associate to any fibration $p : E \rightarrow B$ and any homotopy class of homotopies F between two maps $f_0, f_1 : B' \rightarrow B$ a homotopy class of maps over B' in the form

$$\begin{array}{ccc} f_0^*E & \xrightarrow{\Phi_F} & f_1^*E \\ p_0 \searrow & & \swarrow p_1 \\ & B' & \end{array},$$

relating the two pullback fibrations $p_i : f_i^*E \rightarrow B'$ induced by $f_i : B' \rightarrow B$ for $i = 0, 1$. Complete the proof that this construction defines a functor

$$\Pi(B', B) \rightarrow \mathbf{hTop}_{B'},$$

which associates to each map $f : B' \rightarrow B$ the pullback fibration $f^*E \rightarrow B'$, with the important consequence that the map Φ_F determined by a homotopy is always a homotopy equivalence over B' .

Hint: Consider a family of maps $B' \xrightarrow{f_t} B$ parameterized by $t \in [0, 2]$, which you can think of as a concatenation of a homotopy from f_0 to f_1 with a homotopy from f_1 to f_2 . Defining the induced maps $f_0^*E \rightarrow f_1^*E$ and $f_1^*E \rightarrow f_2^*E$ requires choosing lifts of certain homotopies of maps

$f_0^*E \rightarrow B$ (for $0 \leq t \leq 1$) and $f_1^*E \rightarrow B$ (for $1 \leq t \leq 2$) respectively. Let these choices determine how you can continue the lift of the homotopy of maps $f_0^*E \rightarrow B$ over the interval $1 \leq t \leq 2$, thus defining the induced map $f_0^*E \rightarrow f_2^*E$.

Exercise 4.2. Assume $p : E \rightarrow Y$ is a fibration and $f : X \rightarrow Y$ is a map.

- (a) Show that if $\varphi : X \rightarrow E$ is a map for which the diagram

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & E \\ & \searrow f \quad \sim \quad \swarrow p & \\ & Y & \end{array}$$

commutes up to homotopy, then φ can be replaced with a homotopic map $X \rightarrow E$ that makes the diagram commute on the nose.

- (b) Deduce the basic uniqueness result about fibrations associated to a map $f : X \rightarrow Y$, namely that for any diagram of the form

$$\begin{array}{ccccc} E_0 & \xleftarrow[\sim]{h.e.} & X & \xrightarrow[\sim]{h.e.} & E_1 \\ & \searrow \sim & \downarrow f & \sim \swarrow & \\ & p_0 & Y & p_1 & \end{array}$$

in which $p_0 : E_0 \rightarrow Y$ and $p_1 : E_1 \rightarrow Y$ are both fibrations, the two fibrations are homotopy equivalent over Y .

Exercise 4.3. Suppose \mathcal{C} is a category and $X \xrightarrow{f} Y$ is a morphism in \mathcal{C} with the property that the maps

$$\mathrm{Hom}(Y, X) \xrightarrow{f^*} \mathrm{Hom}(X, X) \quad \text{and} \quad \mathrm{Hom}(Y, Y) \xrightarrow{f^*} \mathrm{Hom}(X, Y)$$

defined via $f^*\varphi := \varphi \circ f$ are both bijections. Prove that f is an isomorphism of \mathcal{C} . Then use this to finish the proof of the theorem stated in lecture that every homotopy equivalence $E \rightarrow E'$ that is also a map over B for two fibrations $E, E' \rightarrow B$ is also a homotopy equivalence over B .

REFERENCES

- [DK01] J. F. Davis and P. Kirk, *Lecture notes in algebraic topology*, Graduate Studies in Mathematics, vol. 35, American Mathematical Society, Providence, RI, 2001.
- [Mac71] S. MacLane, *Categories for the working mathematician*, Graduate Texts in Mathematics, vol. Vol. 5, Springer-Verlag, New York-Berlin, 1971.
- [Mun75] J. R. Munkres, *Topology: a first course*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1975.
- [tD08] T. tom Dieck, *Algebraic topology*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [Wen23] C. Wendl, *Topology I and II* (2023). Notes from the course at HU Berlin, available at <https://www.mathematik.hu-berlin.de/~wendl/Winter2023/Topologie2/lecturenotes.pdf>.